arXiv:hep-ph/9505259v1 8 May 1995

Light Front Quantization

Matthias Burkardt
Institute for Nuclear Theory
University of Washington, Box 351550
Seattle, WA 98195-1550

to appear in:

Advances in Nuclear Physics

Abstract

An introductory overview on Light-Front quantization, with some emphasis on recent achievements, is given. Light-Front quantization is the most promising and physical tool to study deep inelastic scattering on the basis of quark gluon degrees of freedom. The simplified vacuum structure (nontrivial vacuum effects can only appear in zero-mode degrees of freedom) and the physical basis allows for a description of hadrons that stays close to intuition. Recent progress has ben made in understanding the connection between effective LF Hamiltonians and nontrivial vacuum condesates. Discrete Light-Cone Quantization, the transverse lattice and Light-Front Tamm-Dancoff (in combination with renormalization group techniques) are the main tools for exploring LF-Hamiltonians nonperturbatively.

Contents

1 Introduction **3**
- 1.1 Deep Inelastic Scattering . . . 3
- 1.2 Advantages of Light-Front Coordinates . . . 6
- 1.3 Outline . . . 8

2 Canonical Quantization **10**
- 2.1 Quantization in Light-Front Coordinates . . . 10
- 2.2 ε-Coordinates on Finite Light-Front Intervals . . . 13
- 2.3 Examples for Canonical Light-Front Hamiltonians . . . 15
  - 2.3.1 Scalar Fields . . . 16
  - 2.3.2 Fermions with Yukawa Interactions . . . 17
  - 2.3.3 QED and QCD . . . 19

3 The Light-Front Vacuum **23**
- 3.1 The Physical Picture . . . 23
- 3.2 Examples for Zero Modes . . . 25
  - 3.2.1 Constant Scalar Fields . . . 26
  - 3.2.2 Fermionic Zero Modes . . . 27
  - 3.2.3 Gauge Field Zero-Modes . . . 28
  - 3.2.4 Perturbative Zero-Modes . . . 30
- 3.3 Zero Modes and the Vacuum in ε-Coordinates . . . 32
  - 3.3.1 General Considerations . . . 32
  - 3.3.2 A simple Example for the limit $L \to \infty$, $\varepsilon/L \to 0$ . . . 36
- 3.4 Vacuum Condensates and Sum Rules . . . 41

4 Perturbative Renormalization **44**
- 4.1 Scalar Fields . . . 45
- 4.2 Fermions . . . 49
- 4.3 Gauge Theories . . . 53
- 4.4 Summary . . . 54

5 Nonperturbative Calculations **55**
5.1 Discrete Light-Cone Quantization 55
5.2 Functional Integration on a Longitudinal Lattice 59
5.3 Hamiltonian Monte Carlo on a Transverse Lattice 59
5.4 Light-Front Tamm-Dancoff 64

6 Summary, Conclusions and Outlook **69**

A The Dirac Bergmann Formalism **73**

Chapter 1

Introduction

1.1 Deep Inelastic Scattering

More than twenty years after quantum chromodynamics (QCD) was introduced as a microscopic theory of strong interactions, very little is known about its solutions. At least in principle, it should be possible to describe the interaction of nucleons with external probes using quark and gluon degrees of freedom on the basis of QCD. So far, however, the extreme complexity of this theory has slowed any progress in this direction considerably.

Deep inelastic lepton-nucleon scattering (DIS) provides access to quark and gluon degrees of freedom in nucleons and nuclei. In these experiments one shoots high energy leptons (e.g. electrons) at a hadronic target (usually protons or nuclei) and measures the energy and momentum transfer to the target by detecting the final state lepton (Fig. 1.1). The hadronic final state X is not measured (usually the nucleon is destroyed in these reactions and the hadronic final state consists of many particles). Because of the extremely large momentum transfer to the target (typical momentum transfers in DIS experiments are several GeV/c or more), the inclusive cross sections are dominated by single particle response functions along the light-cone. To illustrate this let us use the optical theorem which relates the differential lepton nucleon cross section to the imaginary part of the forward Compton amplitude [Yn 83] (Fig.1.2). One finds

$$\frac{d^2\sigma}{d\Omega dE'} = \frac{\alpha^2}{q^4}\left(\frac{E'}{E}\right) l^{\mu\nu}\frac{\Im T_{\mu\nu}}{2\pi} \tag{1.1}$$

where E, E' are the energies of the initial and final lepton. $q = k - k'$ is the four momentum transfer of the lepton on the target and

Figure 1.1: Inclusive process $e^- + N \to e^{-\prime} + X$, where X is an unidentified hadronic state.

$l_{\mu\nu} = 2k_\mu k'_\nu + 2k_\nu k'_\mu + q^2 g_{\mu\nu}$ is the leptonic tensor. The hadronic tensor

$$T_{\mu\nu}(P,q) = \frac{i}{2M_N} \sum_S \int \frac{d^4x}{2\pi} e^{iq\cdot x} \langle P,S|T\left(J_\mu(x)J_\nu(0)\right)|P,S\rangle \tag{1.2}$$

(S is the spin of the target proton) contains all the information about the parton substructure of the target proton.

In the Bjorken limit ($Q^2 \equiv -q^2 \to \infty$, $P\cdot q \to \infty$, $x_{Bj} = Q^2/2P\cdot q$ fixed), deep inelastic structure functions exhibit Bjorken scaling: up to kinematical coefficients, the hadronic tensor (1.2) depends only on x_{Bj} but no longer on Q^2 (within perturbatively calculable logarithmic corrections). In order to understand this result, it is convenient to introduce *light-front* variables $a_\mp = a^\pm = \left(a^0 \pm a^3\right)/\sqrt{2}$ so that the scalar product reads $a\cdot b = a_+b^+ + a_-b^- - \vec{a}_\perp\vec{b}_\perp = a_+b_- + a_-b_+ - \vec{a}_\perp\vec{b}_\perp$. Furthermore let us choose a frame where $\vec{q}_\perp = 0$. The Bjorken limit corresponds to p^μ and q_- fixed, while $q_+ \to \infty$. Bjorken scaling is equivalent to the statement that the structure functions become independent of q_+ in this limit (again up to trivial kinematic coefficients). In this limit, the integrand in Eq.(1.2) contains the rapidly oscillating factor $\exp(iq_+x^+)$, which kills all contributions to the integral except those where the integrand is singular [Ja 85]. Due to causality, the integrand must vanish for $x^2 = 2x^+x^- - \vec{x}_\perp^2 < 0$ and the current product is singular at $x^+ = 0$, $\vec{x}_\perp = 0$. The leading singularity can be obtained from the operator product expansion by contracting two fermion operators in the product $T\left(J_\mu(x)J_\nu(0)\right) \equiv T\left(\bar\psi(x)\gamma_\mu\psi(x)\bar\psi(0)\gamma_\nu\psi(0)\right)$, yielding a nonlocal term bilinear in the fermion field multiplying a free (asymptotic freedom!)

Figure 1.2: Inclusive lepton nucleon cross section expressed in terms of the imaginary part of the forward Compton amplitude. For $Q^2 = -q^2 \to \infty$ only the 'handbag diagram' (both photons couple to the same quark) survives. The 'crossed diagram' (the two photons couple to different quarks) is suppressed because of wavefunction effects.

fermion propagator from 0 to x which gives rise to the abovementioned singularity structure [CL 84]. The $x^+ = \vec{x}_\perp = 0$ dominance in the integral has two consequences. First it explains Bjorken scaling, because q_+ enters the hadronic tensor only via the term $x^+ q_+$ in the exponent and for $x^+ = 0$ the q_+ dependence drops out. Second, and this is very important for practical calculations, the parton distributions, i.e. the Bjorken scaled structure functions, can be expressed in terms of correlation functions along the *light-front* space direction x^-. For example, for the spin averaged parton distribution one obtains

$$2P_- f(x_{Bj}) = \int \frac{dx^-}{2\pi} \langle P|\bar{\psi}(0)\gamma_-\psi(x^-)|P\rangle \exp(iP_- x^- x_{Bj}), \qquad (1.3)$$

The physical origin of this result can be understood as follows. Consider again the virtual forward Compton amplitude (Fig. 1.2). In principle, the photons in the first and second interaction in Fig. 1.2 can couple to the same as well as to different quarks in the target. However, the hadronic wavefunction can only absorb momenta which are of the order of the QCD-scale ($\Lambda_{QCD} \approx 200 MeV$). Therefore, in the limit of large momentum transfer, only such diagrams survive where the two photons in Fig. 1.2 couple to the same quark. All other diagrams have large momenta flowing through

the wavefunction or they involve extra hard gluon exchanges which results in their suppression at large Q^2. The large momentum transfer is also important because of asymptotic freedom. Since $\alpha_S(Q^2) \sim 1/\log\left(Q^2/\Lambda^2_{QCD}\right)$, the running coupling constant of QCD, goes to zero for large Q^2, all interactions of the struck quark can be neglected and it propagates essentially without interaction between the two photon-vertices. Furthermore, since the momentum transfer is much larger than the masses of the quarks in the target, the struck quarks propagation between becomes ultra-relativistic, i.e. it moves exceedingly close to the light cone $x^2 = 0$. Due to the high-energy nature of the scattering, the relativistic structure function is a LF correlation [Ja 72, Ji 93]. Already at this point it should be clear that LF-coordinates play a distinguished role in the analysis of DIS experiments — a point which will become much more obvious after we have introduced some of the formal ideas of LF quantization.

1.2 Advantages of Light-Front Coordinates

LF quantization is very similar to canonical equal time (ET) quantization [Di 49] (here we closely follow Ref. [Ho 91]). Both are Hamiltonian formulations of field theory, where one specifies the fields on a particular initial surface. The evolution of the fields off the initial surface is determined by the Lagrangian equations of motion. The main difference is the choice of the initial surface, $x^0 = 0$ for ET and $x^+ = 0$ for the LF respectively. In both frameworks states are expanded in terms of fields (and their derivatives) on this surface. Therefore, the same physical state may have very different wavefunctions[1] in the ET and LF approaches because fields at $x^0 = 0$ provide a different basis for expanding a state than fields at $x^+ = 0$. The reason is that the microscopic degrees of freedom — field amplitudes at $x^0 = 0$ versus field amplitudes at $x^+ = 0$ — are in general quite different from each other in the two formalisms.

This has important consequences for the practical calculation of parton distributions (1.3) which are real time response functions in the equal time formalism. [2] In order to evaluate Eq.(1.3) one needs to know not only the ground state wavefunction of the target, but also matrix elements to excited states. In contrast, in the framework of LF quantization, parton distributions are correlation functions at equal LF-time x^+, i.e. *within* the initial surface $x^+ = 0$ and can thus be expressed directly in terms of ground state wavefunctions (As a reminder: ET wavefunctions and LF wavefunctions

[1] By "wavefunction" we mean here the collection of all Fock space amplitudes.

[2] The arguments of $\bar{\psi}$ and ψ in Eq.(1.3) have different time components!

are in general different objects). In the LF framework, parton distributions $f(x_{Bj})$ can be easily calculated and have a very simple physical interpretation as single particle momentum densities, where x_{Bj} measures the fraction of momentum carried by the hadron [3]

$$x_{Bj} = \frac{p_-^{parton}}{P_-^{hadron}}. \tag{1.4}$$

Although DIS is probably the most prominent example for practical applications of LF coordinates, they prove useful in many other places as well. For example, LF coordinates have been used in the context current algebra sum rules in particle physics [FF 65]. Another prominent example is form factors, where moments of the wave function along the LF determine the asymptotic falloff at large momentum transfer [BL 80]. More recently, LF quantization found applications in inclusive decays of heavy quarks [BD+ 92, Bu 92d, CZ+ 95].

From the purely theoretical point of view, various advantages of LF quantization derive from properties of the ten generators of the Poincaré group (translations P^μ, rotations $\vec{L}$ and boosts $\vec{K}$) [Di 49, Ho 91]. Those generators which leave the initial surface invariant ($\vec{P}$ and $\vec{L}$ for ET and P_-, $\vec{P}_\perp$, L_3 and $\vec{K}$ for LF) are "simple" in the sense that they have very simple representations in terms of the fields (typically just sums of single particle operators). The other generators, which include the "Hamiltonians" (P_0, which is conjugate to x^0 in ET and P_+, which is conjugate to the LF-time x^+ in LF quantization) contain interactions among the fields and are typically very complicated. Generators which leave the initial surface invariant are also called *kinematic* generators, while the others are called *dynamic* generators. Obviously it is advantageous to have as many of the ten generators kinematic as possible. There are seven kinematic generators on the LF but only six in ET quantization.

The fact that P_-, the generator of x^- translations, is kinematic (obviously it leaves $x^+ = 0$ invariant!) and positive has striking consequences for the LF vacuum[Ho 91]. For free fields $p^2 = m^2$ implies for the LF energy $p_+ = \left(m^2 + \vec{p}_\perp\right)/2p_-$. Hence positive energy excitations have positive p_-. After the usual reinterpretation of the negative energy states this implies that p_- for a single particle is positive (which makes sense, considering that $p_- = \left(p_0 - p_3\right)/\sqrt{2}$). P_- being kinematic means that it is given by the sum

[3] In DIS with nonrelativistic kinematics (e.g. thermal neutron scattering off liquid ^{4}He) one also observes scaling and the structure functions can be expressed in terms of single particle response functions. However, due to the different kinematics, nonrelativistic structure functions at large momentum transfer are dominated by Fourier transforms of equal time response functions, i.e. ordinary momentum distributions.

of single particle p_-. Combined with the positivity of p_- this implies that the Fock vacuum (no particle excitations) is the unique state with $P_- = 0$. All other states have positive P_-. Hence, even in the presence of interactions, the LF Fock vacuum does not mix with any other state and is therefore an exact eigenstate of the LF Hamiltonian P_+ (which commutes with P_-). If one further assumes parity invariance of the ground state this implies that the Fock vacuum must be the exact ground state of the fully interacting LF quantum field theory. [4] In sharp contrast to other formulations of field theory, the LF-vacuum is trivial! This implies a tremendous technical advantage but also raises the question whether nonperturbative LF-field theory is equivalent to conventional field theory, where nonperturbative effects usually result in a highly nontrivial vacuum structure. This very deep issue will be discussed in more detail in Chapter 3.

Dirac was the first who had the idea to formulate field theory in LF-coordinates [Di 49].[5] In this remarkable work (almost 20 years before scaling was discovered in deep inelastic lepton nucleon scattering !) he has shown that it should in principle be possible to formulate a consistent quantum theory on the LF. This work laid the basis for all further developments, but left many details open. The main issues are the structure of the vacuum, renormalization and practical algorithms for solution.

1.3 Outline

There are many similarities between the formal steps in ET quantization and LF quantization. In Chapter 2 we will explain the basic steps in constructing LF Hamiltonians and give examples for scalar fields, fermions and gauge fields. The vacuum on the LF is very controversial. On the one hand simple kinematical arguments seem to show that in LF field theory the vacuum of interacting field theories is the same as the free field theory vacuum (all interactions turned off). In QCD we know that chiral symmetry is spontaneously broken. It is up to now unclear whether a LF Hamiltonian, with its trivial vacuum, is capable of describing this physics. We will elaborate on this point in Chapter 3. Renormalization is an issue because the LF-approach to field theory is not manifestly covariant. Thus UV-divergences (which occur on the LF as they do everywhere in quantum field theory) are not necessarily

[4] Practical calculations show that typical LF Hamiltonians are either unbounded from below or their ground state is indeed the Fock vacuum.

[5] Later, a similar framework was developed independently on the basis of a Lorentz frame ("the infinite momentum frame") that moves with $v \to c$ [FF 65, Su 68, We 69, KS 70, BK+ 71, BR+ 73].

the same for all Lorentz components of a particular operator under consideration. Clearly this requires a more complex counterterm structure to render the theory finite and to restore Lorentz invariance for physical observables (see Chapter 4). Despite certain technical simplifications, field theory on the LF is *a priori* still an enormously complex many body problem. In particular in QCD one knows from DIS experiments that the nucleon consists not only of the three valence quarks, but that sea quark pairs and gluons are a significant, if not dominant, component of the nucleon's LF wavefunction, i.e. one should not expect that the LF wavefunctions of ground state hadrons in QCD are simple. Recent attempts to cast LF bound state problems into a form that can be solved on a computer will be described in Chapter 5.

Chapter 2

Canonical Quantization

2.1 Quantization in Light-Front Coordinates

In this chapter, the formal steps for quantization on the light-front are presented. For pedagogical reasons this will be done by comparing with conventional quantization (with x^0 as "time"). On the one hand this shows that the basic steps in the quantization procedure in LF and in ET formalism are in fact very similar. More importantly, however, we will thus be able to highlight the essential differences between these two approaches to quantum field theory more easily.

In the context of canonical quantization one usually starts from the action

$$S = \int d^4x \mathcal{L}. \tag{2.1}$$

($\mathcal{L} = \mathcal{L}(\phi, \partial_\mu \phi)$) After selecting a time direction τ [1] one forms the momenta which are canonically conjugate to ϕ

$$\Pi(x) = \frac{\delta \mathcal{L}}{\delta \partial_\tau \phi} \tag{2.2}$$

and postulates canonical commutation relations between fields and corresponding momenta at equal "time" τ (Table 2.1). [2]

In the next step one constructs the Hamilton operator and the other components of the momentum vector. Thus one has completely specified the

[1] Here τ may stand for ordinary time x^0 as well as for LF time $x^+ = \left(x^0 + x^3\right)/\sqrt{2}$ or any other (not space-like) direction.

[2] The canonical quantization procedure in the ET formulation can for example be found in Ref. [BD 65]. The rules for canonical LF-quantization have been taken from Refs. [CR+ 73, CY 72].

normal coordinates	light-front
coordinates	
x^0 time x^1, x^2, x^3 space	$x^+ = \frac{x^0 + x^3}{\sqrt{2}}$ time $x^- = \frac{x^0 - x^3}{\sqrt{2}}, x^1, x^2$ space
scalar product	
$a \cdot b = a^0 b^0 - a^1 b^1 - a^2 b^2 - a^3 b^3 = a^0 b^0 - \vec{a}\vec{b}$	$a \cdot b = a^+ b^- + a^- b^+ - a^1 b^1 - a^2 b^2 = a^+ b^- + a^- b^+ - \vec{a}_\perp \vec{b}_\perp$
Lagrangian density	
$\mathcal{L} = \frac{1}{2}(\partial_0 \phi)^2 - \frac{1}{2}\left(\vec{\nabla}\,\phi\right)^2 - V(\phi)$	$\mathcal{L} = \partial_+ \phi \partial_- \phi - \frac{1}{2}\left(\vec{\nabla}_\perp\,\phi\right)^2 - V(\phi)$
conjugate momenta	
$\pi = \frac{\delta \mathcal{L}}{\delta \partial_0 \varphi} = \partial_0 \varphi$	$\pi = \frac{\delta \mathcal{L}}{\delta \partial_+ \varphi} = \partial_- \varphi$
canonical commutation relations	
$[\pi(\vec{x}, t), \varphi(\vec{y}, t)] = -i\delta^3(\vec{x} - \vec{y})$	$[\pi(x^-, x_\perp, x^+), \varphi(y^-, y_\perp, x^+)] = -\frac{i}{2}\delta(x^- - y^-)\delta^2(\vec{x}_\perp - \vec{y}_\perp)$
Hamilton operator	
$P^0 = \int d^3x\, \mathcal{H}(\varphi, \pi)$ $\mathcal{H} = \pi \partial_0 \varphi - \mathcal{L}$	$P_+ = \int dx^- \int d^2x_\perp\, \mathcal{H}(\varphi, \pi)$ $\mathcal{H} = \pi \partial_+ \varphi - \mathcal{L}$
momentum operator	
$\vec{P} = \int d^3x\, \pi \vec{\bigtriangledown} \varphi$	$P_- = \int dx^- d^2x_\perp\, \pi \partial_- \varphi$ $\vec{P}_\perp = \int dx^- d^2x_\perp\, \pi \vec{\partial}_\perp \varphi$
eigenvalue equation	
$P^0 \|\psi_n> = E_n \|\psi_n>$ $\vec{P}$ fixed	$P_+ \|\psi_n> = P_{+n} \|\psi_n>$ $P_-, \vec{P}_\perp$ fixed
hadron masses	
$M_n^2 = E_n^2 - \vec{P}^2$	$M_n^2 = 2P_{+n}P_- - \vec{P}_\perp^2$

Table 2.1: canonical quantization in ordinary coordinates and on the light-front

dynamics and can start solving the equations of motion. Typically, one either makes some variational ansatz or a Fock space expansion. In the latter approach one writes the hadron wave function as a sum over components with a fixed number of elementary quanta (for example in QCD: $q\bar{q}$, $q\bar{q}q\bar{q}$, $q\bar{q}g$, e.t.c.). The expansion coefficients, i.e. the wavefunctions for the corresponding Fock space sector are used as variational parameters. They are determined by making the expectation value of the energy stationary with respect to variations in the wavefunction. Typically the variation is done for fixed momentum.[3] This whole procedure results in coupled integral equations for the Fock space components. In general they have to be solved numerically. In practical calculations, since one cannot include infinitely many Fock components, one has to introduce some *ad hoc* cutoff in the Fock space. Thus it is very important to demonstrate that physical observables do not depend on how many Fock components are included.

Until one selects the canonically conjugate momenta and postulates equal τ commutation relations, i.e. at the level of the classical Lagrangian, the transition from ET to the LF consists of a mere rewriting. After quantization, the independent degrees of freedom consist of the fields and their conjugate momenta on the initial surface ($x^0 = 0$ for ET and $x^+ = 0$ for LF). Thus different degrees of freedom are employed to expand physical states in the ET and in the LF approach. Of course, after solving the equations of motion, physical observables must not depend on the choice of quantization plane. However, it may turn out that one approach is more efficient (e.g. faster numerical convergence) than the other or more elegant and more easy to interpret physically. In general, this will of course depend on the details of the interaction. An extreme example is $\mathrm{QCD}_{1+1}(N_C \to \infty)$. In the ET approach [BG 78, Li 86, HN+ 88] one first has to solve coupled, nonlinear integral equations with a singular kernel to obtain the Hartree-Fock solution for the vacuum. Then, in order to calculate meson masses, one has to solve the two body equation in this background, which amounts to solving another set of coupled (linear) integral equations with singular kernel. In the LF-approach [Ho 74, Ei 76] all one has to do is solve one linear integral equation with singular kernel. The numerical results for the meson spectrum are in extremely good agreement between the two approaches, but numerically the LF calculation is more than one order of magnitude faster! In this case the simplification arises because the LF-vacuum is trivial — a point which will be elaborated in more detail below as well as in Chapter 3.

Which approach is preferable may, however, also depend on the observ-

[3] On the LF this is very important because $P_+ \propto 1/P_-$, i.e. unrestricted variation (P_- allowed to vary) results in $P_- \to \infty$.

ables in which one is interested. The most prominent example is deep inelastic scattering. As discussed in the introduction, parton distributions are much more easily accessible on the LF than in usual coordinates.

2.2 ε-Coordinates on Finite Light-Front Intervals

One issue one may be worried about is the question of equivalence between the LF approach to field theories and other approaches. On the LF one imposes commutation relations at equal LF-time, i.e. between two space-time points that are connected by a light-like distance. Thus it is *a priori* not clear whether the initial value problem with initial conditions on a null plane is well defined [Ro 70, RM 92] and whether there arise any conflicts with causality on the LF. The situation becomes particularly worrisome when one introduces a "box" in the longitudinal x^- direction (to keep IR-singularities under control) and imposes periodic or quasiperiodic boundary conditions at the ends of the box — i.e. one imposes boundary conditions between points that may be causally related. One way to address this issue in a well defined way is to define the LF via a limiting procedure by starting from a spacelike quantization surface and carefully rotating this surface until one has 'reached' the LF (note: although there are some similarities, this should not be confused with a Lorentz boost to infinite momentum [We 69, BR+ 73]).

In order to be able to control infrared singularities, let us formulate the dynamics on a finite LF interval with extension L in the x^- direction. [4] On a finite interval, boundary conditions have to be specified, e.g. $\phi(x^- + L, x^+) = \phi(x^-, x^+)$. However, if one is working on the LF, imposing boundary conditions means relating fields at points that are separated by a light-like distance — obviously one may run into trouble with causality at this point. To avoid this dilemma, Lenz et al. [LT 91] have introduced ε-coordinates which are defined as follows, [5]

$$\begin{aligned} x_\varepsilon^- &= x^- \\ x_\varepsilon^+ &= x^+ + \frac{\varepsilon}{L} x^-. \end{aligned} \tag{2.3}$$

Now points at opposite ends of the interval (with coordinates $(x_\varepsilon^- + L, x_\varepsilon^+)$ and $(x_\varepsilon^-, x_\varepsilon^+)$ are separated by a spacelike distance $ds^2 = -2\varepsilon L$ and no conflict

[4] To simplify the notation, only 1+1 dimensional examples will be discussed in this section.

[5] See also Ref.[PF 89]. A slightly different approach, where both x^+ and x^- are rotated away from the light-cone, has been studied in Ref.[Ho 92].

with causality arises from imposing boundary conditions. In ε-coordinates the scalar product is given by

$$A \cdot B = A_+ B_- + A_- B_+ + 2\frac{\varepsilon}{L} A_+ B_+ \tag{2.4}$$

and thus the Lagrangian density (for the rest of this section, the subscript ε will be dropped to simplify the notation) for ϕ^4_{1+1} reads

$$\mathcal{L} = \partial_+\phi \left(\partial_-\phi + \frac{\varepsilon}{L}\partial_+\phi \right) - \frac{m^2}{2}\phi^2 - \frac{\lambda}{4!}\phi^4. \tag{2.5}$$

Since $\mathcal{L}$ is quadratic in $\partial_+\phi$, quantization in ε-coordinates is straightforward (as in usual coordinates). One finds [LT 91]

$$\Pi = \frac{\delta \mathcal{L}}{\delta \partial_+ \phi} = \partial_-\phi + \frac{2\varepsilon}{L}\partial_+\phi \tag{2.6}$$

$$\left[\Pi(x^-, x^+), \phi(y^-, x^+) \right] = -i\delta(x^- - y^-) \tag{2.7}$$

and

$$H = \int dx^- \frac{L}{4\varepsilon} \left(\Pi - \partial_-\phi \right)^2 + \frac{m^2}{2}\phi^2 + \frac{\lambda}{4!}\phi^4. \tag{2.8}$$

In these coordinates, the free dispersion relation ($\lambda = 0$) is given by

$$p_+(n) = \frac{L}{2\varepsilon} \left(-p_-(n) \pm \sqrt{p_-(n)^2 + \frac{2\varepsilon}{L} m^2} \right), \tag{2.9}$$

where $p_-(n) = \frac{2\pi n}{L}$ as usual in a box with periodic boundary conditions. Later, we will also need the normal mode expansion of the fields with periodic boundary conditions

$$\begin{aligned} \phi(x^-) &= \sum_n \frac{1}{2\sqrt{\omega_n}} \left[a_n e^{-ip_-(n)x^-} + a_n^\dagger e^{ip_-(n)x^-} \right] \\ \Pi(x^-) &= \sum_n \frac{-i\sqrt{\omega_n}}{L} \left[a_n e^{-ip_-(n)x^-} - a_n^\dagger e^{ip_-(n)x^-} \right], \end{aligned} \tag{2.10}$$

where $\omega_n = L\sqrt{p_-(n)^2 + \frac{2\varepsilon}{L}m^2}$ and the a, $a^\dagger$ satisfy the usual commutation relations, e.g

$$\left[a_m, a_n^\dagger \right] = \delta_{m,n}. \tag{2.11}$$

The most significant difference between the dispersion relation in ε-coordinates (2.9) and the dispersion relation on the LF ($p_+ = \frac{m^2}{2p_-}$) is the appearance of two solutions of p_+ for each p_- in ε-coordinates, while the

Figure 2.1: Free dispersion relation in ε-coordinates versus the dispersion relation in the LF limit.

dispersion relation on the LF yields just one solution for each p_- (Figure 2.1). For positive energy ($p_+ > 0$) modes, the LF momentum p_- is positive whereas the momentum p_- in ε-coordinates can be both positive and negative. This has importance consequences for the vacuum structure which will be discussed in Chapter 3.

In the limit $\frac{\varepsilon}{L} \to 0$ (L fixed) the LF is recovered:

$$\Pi \overset{\frac{\varepsilon}{L}\to 0}{\longrightarrow} \partial_- \phi \tag{2.12}$$

$$H \overset{\frac{\varepsilon}{L}\to 0}{\longrightarrow} \int dx^- \frac{m^2}{2}\phi^2 + \frac{\lambda}{4!}\phi^4. \tag{2.13}$$

For all nonzero ε, the relation between the momenta and the fields (2.6) contains the time derivative of the fields and the fields are quantized as usual (2.7). However, for $\varepsilon = 0$, Eq.(2.6) becomes a constraint equation, and the Dirac-Bergmann algorithm (see Appendix A) yields $[\partial_- \phi(x^-, x^+), \phi(y^-, x^+)] = \frac{i}{2}\delta(x^- - y^-)$.

It should be noted, that the order of limits does matter, i.e. it is important whether one takes the LF limit ($\frac{\varepsilon}{L} \to 0$) first or the continuum limit ($L \to \infty$). This will be discussed in detail in Chapter 3.

2.3 Examples for Canonical Light-Front Hamiltonians

2.3.1 Scalar Fields

Self-interacting Scalar fields in the LF framework have been discussed in Refs.[CR+ 73, CY 72]. In order to keep the discussion as general as possible, we will work in $D_\perp$ transverse dimensions, where $D_\perp = 0, 1, 2$. For a polynomial interaction [6],

$$\mathcal{L} = \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - \frac{m^2}{2}\phi^2 - \mathcal{L}^{int}, \tag{2.14}$$

where $\mathcal{L}^{int}$ is a polynomial in ϕ, the momenta conjugate to ϕ are

$$\Pi = \partial_-\phi \tag{2.15}$$

with commutation relations

$$\Big[\Pi(x^-, \vec{x}_\perp, x^+), \phi(y^-, \vec{y}_\perp, x^+)\Big] = -\frac{i}{2}\delta(x^- - y^-)\delta(\vec{x}_\perp - \vec{y}_\perp). \tag{2.16}$$

Note that this implies nonlocal commutation relations for the field ϕ, e.g.

$$\Big[\phi(x^-, \vec{x}_\perp, x^+), \phi(y^-, \vec{y}_\perp, x^+)\Big] = -\frac{i}{4}\varepsilon(x^- - y^-)\delta(\vec{x}_\perp - \vec{y}_\perp), \tag{2.17}$$

where $\varepsilon(x) = 1$ for $x > 0$ and $\varepsilon(x) = -1$ for $x < 0$. The Hamiltonian density ($P_+ = \int dx^- d^{D_\perp}x_\perp \mathcal{H}$) is obtained from Eq.(2.14) via a Legendre transformation

$$\begin{aligned} \mathcal{H} &= \Pi\partial_+\phi - \mathcal{L} \\ &= \frac{1}{2}\left(\vec{\nabla}_\perp\phi\right)^2 + \frac{m^2}{2}\phi^2 + \mathcal{L}^{int}. \end{aligned} \tag{2.18}$$

The commutation relations (2.16) are easily satisfied if we make a mode expansion

$$\phi(x) = \int_0^\infty \frac{dk_-}{\sqrt{4\pi k_-}} \int \frac{d^{D_\perp}k_\perp}{(2\pi)^{D_\perp/2}} \left[a_{k_-\vec{k}_\perp} e^{-ikx} + a^\dagger_{k_-\vec{k}_\perp} e^{ikx}\right] \tag{2.19}$$

where $a_{k_-\vec{k}_\perp}$, $a^\dagger_{k_-\vec{k}_\perp}$ satisfy the usual boson commutation relations, e.g.

$$\left[a_{k_-\vec{k}_\perp}, a^\dagger_{q_-\vec{q}_\perp}\right] = \delta(k_- - q_-)\delta(\vec{k}_\perp - \vec{q}_\perp). \tag{2.20}$$

[6] In 3+1 dimensions, renormalizability restricts the interaction to 4th order polynomials, but in $2+1$ or $1+1$ dimensions higher order polynomials are conceivable (6th order and ∞ order respectively).

Longitudinal and transverse momentum operators contain no interaction terms

$$\begin{aligned} P_- &= \int dx^- d^{D_\perp} x_\perp \Pi \partial_- \phi = \int_0^\infty dk_- \int d^{D_\perp} k_\perp k_- a^\dagger_{k_- \vec{k}_\perp} a_{k_- \vec{k}_\perp} \\ \vec{P}_\perp &= \int dx^- d^{D_\perp} x_\perp \Pi \vec{\nabla}_\perp \phi = \int_0^\infty dk_- \int d^{D_\perp} k_\perp \vec{k}_\perp a^\dagger_{k_- \vec{k}_\perp} a_{k_- \vec{k}_\perp} \end{aligned} \quad (2.21)$$

where normal ordering terms have been dropped. Most of the numerical studies of self-interacting scalar fields have been done in $1+1$ dimensions [HV 87, Bu 93] using discrete light-cone quantization (Section 5.1). A more recent work employs Monte Carlo techniques to solve ϕ^4-theory in $2+1$ dimensions [Bu 94a].

Complex scalar fields can always be reduced to real scalar fields by working in a Cartesian basis $\Phi = \frac{1}{\sqrt{2}}(\phi_1 + i\phi_2)$ and thus need not be discussed here.

2.3.2 Fermions with Yukawa Interactions

To keep the discussion as general as possible we assume an interaction of the form $\bar{\psi}\Gamma\psi\phi$, where ϕ is either scalar or pseudoscalar and Γ is either 1 or $i\gamma_5$

$$\mathcal{L} = \bar{\psi}\left(i \not{\partial} - M - g\Gamma\phi\right)\psi + \frac{1}{2}\left(\partial_\mu\phi\partial^\mu\phi - m^2\phi^2\right). \quad (2.22)$$

One novel feature compared to normal coordinates and compared to self-interacting scalar fields on the LF is the fact that not all components of ψ are independent dynamical degrees of freedom. To see this, let us introduce projection matrices $\mathcal{P}^{(\pm)} = \frac{1}{2}\gamma^\mp\gamma^\pm$ where $\gamma^\pm = (\gamma^0 \pm \gamma^3)/\sqrt{2}$. Note that $\gamma^+\gamma^+ = \gamma^-\gamma^- = 0$ implies $\mathcal{P}^{(+)}\mathcal{P}^{(-)} = \mathcal{P}^{(-)}\mathcal{P}^{(+)} = 0$. These projection matrices can be used to decompose the fermion spinors into dynamical and non-dynamical components $\psi = \psi_{(+)} + \psi_{(-)}$,where $\psi_{(\pm)} \equiv \mathcal{P}^{(\pm)}\psi$. The Lagrangian does not contain a LF-time derivative (∂_+) of $\psi_{(-)}$

$$\begin{aligned} \mathcal{L} &= \sqrt{2}\psi^\dagger_{(+)} i\partial_+ \psi_{(+)} + \sqrt{2}\psi^\dagger_{(-)} i\partial_- \psi_{(-)} - \psi^\dagger_{(+)}\left(i\vec{\alpha}_\perp\vec{\partial}_\perp + \gamma^0\mathcal{M}\right)\psi_{(-)} \\ &\quad -\psi^\dagger_{(-)}\left(i\vec{\alpha}_\perp\vec{\partial}_\perp + \gamma^0\mathcal{M}\right)\psi_{(+)} + \frac{1}{2}\left(\partial_\mu\phi\partial^\mu\phi - m^2\phi^2\right), \end{aligned} \quad (2.23)$$

where $\vec{\alpha}_\perp = \gamma^0\vec{\gamma}_\perp$ and $\mathcal{M}(x) = M + g\Gamma\phi(x)$. Thus the Euler-Lagrange equation for $\psi_{(-)}$ is a constraint equation

$$\sqrt{2}i\partial_-\psi_{(-)} = \left(i\vec{\alpha}_\perp\vec{\partial}_\perp + \gamma^0\mathcal{M}\right)\psi_{(+)}. \quad (2.24)$$

It is therefore necessary to eliminate the dependent degrees of freedom ($\psi_{(-)}$) before quantizing the theory. Here we proceed by solving Eq.(2.24) and inserting the solution back in the Lagrangian (2.23), yielding [CY 72] [7]

$$\begin{aligned}\mathcal{L}_{(+)} &= \sqrt{2}\psi^{\dagger}_{(+)}\partial_{+}\psi_{(+)} + \frac{1}{2}\left(\partial_{\mu}\phi\partial^{\mu}\phi - m^2\phi^2\right) \\ &\quad -\frac{1}{\sqrt{2}}\psi^{\dagger}_{(+)}\left(i\vec{\alpha}_{\perp}\vec{\partial}_{\perp} + \gamma^0\mathcal{M}\right)\frac{1}{i\partial_{-}}\left(i\vec{\alpha}_{\perp}\vec{\partial}_{\perp} + \gamma^0\mathcal{M}\right)\psi_{(+)}. \quad (2.25)\end{aligned}$$

The ambiguities associated with the inversion of the differential operator will be discussed in Section 3.2.2. Here we just define

$$\left(\frac{1}{\partial_{-}}f\right)(x^{-},\vec{x}_{\perp}) = \frac{1}{2}\int_{-\infty}^{\infty} dy^{-}\varepsilon(x^{-}-y^{-})f(y^{-},\vec{x}_{\perp}). \quad (2.26)$$

The rest of the quantization procedure is now straightforward. The Hamiltonian is given by

$$P^{-} = \int dx^{-}d^{D_{\perp}}x_{\perp}\mathcal{H} \quad (2.27)$$

where

$$\begin{aligned}\mathcal{H} &= \frac{1}{\sqrt{2}}\psi^{\dagger}_{(+)}\left(i\vec{\alpha}_{\perp}\vec{\partial}_{\perp} + \gamma^0\mathcal{M}\right)\frac{1}{i\partial_{-}}\left(i\vec{\alpha}_{\perp}\vec{\partial}_{\perp} + \gamma^0\mathcal{M}\right)\psi_{(+)} \\ &\quad +\frac{1}{2}\left[\left(\vec{\nabla}_{\perp}\phi\right)^2 + m^2\phi^2\right] \quad (2.28)\end{aligned}$$

Note that Eq.(2.28) contains four-point interactions of the form $\psi^{\dagger}_{(+)}\phi\left(i\partial_{-}\right)^{-1}\phi\psi_{(+)}$ which were not present in the original Lagrangian (2.22). Note also, that the fermion mass M enters the Hamiltonian density (2.28) in two different places: in the kinetic term for the dynamical fermion field, $\mathcal{H}_{kin} \propto M^2\psi^{\dagger}_{(+)}\left(i\partial_{-}\right)^{-1}\psi_{(+)}$, as well as in the three point vertex, $Mg\psi^{\dagger}_{(+)}\left(i\partial_{-}\right)^{-1}\Gamma\phi\psi_{(+)} + h.c.$. In Chapter 4 we will find that in general these two masses are renormalized differently.

The scalar field ϕ is quantized as in Section (2.3.1). For the fermions, one imposes anti-commutation relations only for the independent component $\psi_{(+)}$ [CY 72]

$$\left\{\psi_{(+)}(x),\psi^{\dagger}_{(+)}(y)\right\}_{x^{+}=y^{+}} = \frac{1}{\sqrt{2}}\mathcal{P}^{(+)}\delta(x^{-}-y^{-})\delta(\vec{x}_{\perp}-\vec{y}_{\perp}) \quad (2.29)$$

with $\left\{\psi^{\dagger}_{(+)}(x),\psi^{\dagger}_{(+)}(y)\right\}_{x^{+}=y^{+}}$ and $\left\{\psi_{(+)}(x),\psi_{(+)}(y)\right\}_{x^{+}=y^{+}}$ both vanishing.

[7] Another option would be to use Dirac-Bergmann quantization (Appendix A). Up to possible differences in the zero-mode sector, the result is the same.

For practical calculations it is very useful to make a mode expansion. Let $u(p,s)$, $v(p,s)$ be the usual particle and antiparticle spinors, satisfying $(\not{p} - M)\, u(p,s) = 0$ and $(\not{p} + M)\, v(p,s) = 0$, where s labels the spin. The normalization is fixed such that

$$\begin{aligned} \sqrt{2} u^\dagger_{(+)}(p,s') u_{(+)}(p,s) &= \bar{u}(p,s')\gamma_- u(p,s) = 2p_- \delta_{ss'} \\ \sqrt{2} v^\dagger_{(+)}(p,s') v_{(+)}(p,s) &= \bar{v}(p,s')\gamma_- v(p,s) = 2p_- \delta_{ss'}. \end{aligned} \tag{2.30}$$

For $\psi_{(+)}$ we make a plane wave ansatz

$$\psi_{(+)}(x) = \int \frac{d^{D_\perp} p_\perp}{(2\pi)^{D_\perp/2}} \int_0^\infty \frac{dp_-}{\sqrt{4\pi p_-}} \sum_s \left[b_{p_- \vec{p}_\perp s} u_{(+)}(p,s) e^{-ipx} + d^\dagger_{p_- \vec{p}_\perp s} v_{(+)}(p,s) e^{ipx} \right]. \tag{2.31}$$

One can easily verify that $\psi_{(+)}$ in Eq.(2.31) satisfies the anti-commutation relations above (2.29), provided

$$\begin{aligned} \left\{ b^\dagger_{p_- \vec{p}_\perp r}, b_{q_- \vec{q}_\perp s} \right\} &= \delta(p_- - q_-)\delta(\vec{p}_\perp - \vec{q}_\perp)\delta_{rs} \\ \left\{ d^\dagger_{p_- \vec{p}_\perp r}, d_{q_- \vec{q}_\perp s} \right\} &= \delta(p_- - q_-)\delta(\vec{p}_\perp - \vec{q}_\perp)\delta_{rs} \end{aligned} \tag{2.32}$$

with all other anti-commutators vanishing.

Nonperturbative numerical works on the LF Hamiltonian with Yukawa interactions (2.28) have been restricted to DLCQ calculations in 1+1 dimensions [PB 85] as well as to 1+1 dimensional [HP 91] 3+1 dimensional [Wo 93] calculations which use Tamm-Dancoff truncations to fermion and at most two bosons or fermion, antifermion and at most one boson.

2.3.3 QED and QCD

Before one can canonically quantize a gauge field theory, one must fix the gauge — otherwise one has to deal with the infinite degeneracy associated with the gauge symmetry.

In the context of LF quantization one usually picks the LF-gauge $A^+ = A_- = 0$. There are several (related) reasons for this choice. In QED, the constraint equation for the "bad" spinor component reads

$$\sqrt{2} i D_- \psi_{(-)} \equiv \sqrt{2}\,(i\partial_- - eA_-)\,\psi_{(-)} = \left(i\vec{\alpha}_\perp \vec{D}_\perp + \gamma^0 M \right) \psi_{(+)}. \tag{2.33}$$

The solution to this constraint equation,

$$\psi_{(-)} = \frac{1}{\sqrt{2}}\,(i\partial_- - eA_-)^{-1} \left(i\vec{\alpha}_\perp \vec{D}_\perp + \gamma^0 M \right) \psi_{(+)}, \tag{2.34}$$

contains A_- in the denominator and, unless one chooses $A_- = 0$ gauge, one thus obtains terms which have A_- in the denominator in the LF-Hamiltonian. In other words, in any gauge other than the LF-gauge the canonical LF-Hamiltonian always contains all powers of A_- (after expanding the geometric series) appearing in the interactions.

In QCD one faces the additional problem that $\vec{\Pi}_\perp$, the momentum conjugate to $\vec{A}_\perp$, satisfies a nonlinear constraint equation if $A_- \neq 0$ (Section 3.2.3). Another reason to pick LF gauge is that $A_- = 0$ is invariant under the kinematic Lorentz symmetries of the LF, i.e. under all transformations that leave the plane $x^+ = 0$ invariant. It is for these reasons that the LF gauge has been commonly used for canonical LF quantization of gauge field theories, and will also be used here — despite all the difficulties which are inherent to the LF and axial gauges [Ma 83, BD+ 85, Wu 77, LN+ 94a, LN+ 94b, LN+ 94c].

Even after fixing the gauge, not all degrees of freedom are dynamical (similar to $\psi_{(-)}$, their time derivative does not enter the Lagrangian). Before we can proceed with the canonical quantization we first have to eliminate these dependent variables by solving those equations of motion which are constraint equations. For $\psi_{(-)}$ we use Eq.(2.34) (note: $A_- = 0$) and proceed similar to the example of the Yukawa theory. Since the time derivative of A_+ does not enter the Lagrangian, A_+ has to be eliminated as well, by solving its constraint equation (obtained by varying the Lagrangian density with respect to A_+)

$$\partial_-^2 A_+ = \partial_- \vec{\nabla}_\perp \vec{A}_\perp - j^+, \tag{2.35}$$

where $j^+ = \sqrt{2} e \psi^\dagger_{(+)} \psi_{(+)}$, in QED and

$$\partial_-^2 A_{+a} = \partial_- \vec{\nabla}_\perp \vec{A}_{\perp a} + g f^{abc} \vec{A}_{\perp b} \partial_- \vec{A}_{\perp c} - j^+_a, \tag{2.36}$$

where $j^+_a = \sqrt{2} g \psi^\dagger_{(+)\alpha} \psi_{(+)\beta} \frac{\lambda_a^{\alpha\beta}}{2}$, in QCD. After inserting A_+ back into the Lagrangian one can proceed with the quantization as usual. One finds

$$\begin{aligned} \mathcal{H}_{QED} &= \frac{1}{\sqrt{2}} \psi^\dagger_{(+)} \left(i\vec{\alpha}_\perp \vec{D}_\perp + \gamma^0 M \right) \frac{1}{i\partial_-} \left(i\vec{\alpha}_\perp \vec{D}_\perp + \gamma^0 M \right) \psi_{(+)} \\ &\quad - \frac{1}{2} \left(\partial_- \vec{\nabla}_\perp \vec{A}_\perp - j^+ \right) \frac{1}{\partial_-^2} \left(\partial_- \vec{\nabla}_\perp \vec{A}_\perp - j^+ \right), \end{aligned} \tag{2.37}$$

and

$$\begin{aligned} \mathcal{H}_{QCD} &= \frac{1}{\sqrt{2}} \psi^\dagger_{(+)} \left(i\vec{\alpha}_\perp \vec{D}_\perp + \gamma^0 M \right) \frac{1}{i\partial_-} \left(i\vec{\alpha}_\perp \vec{D}_\perp + \gamma^0 M \right) \psi_{(+)} \\ &\quad - \frac{1}{2} \left(\vec{D}_\perp \partial_- \vec{A}_{\perp a} - j^+_a \right) \frac{1}{\partial_-^2} \left(\vec{D}_\perp \partial_- \vec{A}_{\perp a} - j^+_a \right) \end{aligned} \tag{2.38}$$

where $\vec{D}_\perp \partial_- \vec{A}_{\perp a} = \vec{\nabla}_\perp \partial_- \vec{A}_{\perp a} + g f^{abc} \vec{A}_{\perp b} \partial_- \vec{A}_{\perp c}$. The commutation relations are similar to the ones in Yukawa theory

$$\begin{aligned}
\text{QED:} \qquad & \left\{ \psi_{(+)}(x), \psi^\dagger_{(+)}(y) \right\}_{x^+=y^+} = \frac{1}{\sqrt{2}} \mathcal{P}^{(+)} \delta(x^- - y^-) \delta(\vec{x}_\perp - \vec{y}_\perp) \\
& \left[\partial_- A_{\perp i}(x), A_{\perp j}(y) \right]_{x^+=y^+} = -\frac{i}{2} \delta(x^- - y^-) \delta(\vec{x}_\perp - \vec{y}_\perp) \delta_{ij} \\
\text{QCD:} \qquad & \left\{ \psi_{(+)\alpha}(x), \psi^\dagger_{(+)\beta}(y) \right\}_{x^+=y^+} = \frac{1}{\sqrt{2}} \mathcal{P}^{(+)} \delta(x^- - y^-) \delta(\vec{x}_\perp - \vec{y}_\perp) \delta_{\alpha\beta} \\
& \left[\partial_- A_{\perp ai}(x), A_{\perp bj}(y) \right]_{x^+=y^+} = -\frac{i}{2} \delta(x^- - y^-) \delta(\vec{x}_\perp - \vec{y}_\perp) \delta_{ij} \delta_{ab}.
\end{aligned} \tag{2.39}$$

Similar to the approach to scalar field theories and Yukawa theories, one may now attempt to solve the above Hamiltonians by making a mode expansion and using matrix diagonalization. In $1+1$ dimension this method was very successful [EP+ 87, EP 89, HP+ 90, Bu 89a, Bu 89b, BB 91]. In $3+1$ dimensions, this approach suffers from a fundamental problem [8]: charged particles are subject to a linear, confining interaction — which is present even in $\mathcal{H}_{QED}$. For gauge invariant amplitudes (all intermediate states included, which contribute to a given order of the coupling) this linear potential is canceled by infrared singular couplings of charges to the $\perp$-components of the gauge field. However, in most practical calculations, drastic truncations of the Fock space are used to keep the dimension of the Hamiltonian matrix within practical limits [TB 91, KP 92]. This approximation results in incomplete cancelations of IR singularities and IR divergences result. Partly responsible for this disaster is an improper treatment of zero-modes and incomplete gauge fixing. If one integrates the Maxwell equation for $F^{\mu+}$ over x^- one finds [KP 93]

$$-\vec{\partial}_\perp^2 \int dx^- A_-(x^+, x^-, x_\perp) = \int dx^- j^+(x^+, x^-, x_\perp), \tag{2.40}$$

i.e. in general, when $\int dx^- j^+(x^+, x^-, x_\perp) \neq 0$, the "gauge" $A_- = 0$ is inconsistent with the equations of motion. On a finite interval, with periodic boundary condition, this becomes clearer because then a Wilson loop "around the torus", $\exp\left(ie \oint dx^- A_-(x^+, x^-, x_\perp)\right)$, is a gauge invariant quantity. The closest one can get to the LF gauge is $\partial_- A_- = 0$. In this gauge one can now investigate the problem of incomplete gauge fixing. The gauge $\partial_- A_- = 0$ still leaves the freedom of x^--independent gauge transformations $A_\mu \to A'_\mu = A_\mu + \partial_\mu \chi$ where $\partial_-^2 \chi = 0$ (or $\partial_- \chi = 0$ if we restrict ourselves

[8] Besides numerical difficulties which will be discussed in Section 5.1.

to periodic χ) [KP 93]. In such an incompletely gauge fixed situations, not all degrees of freedom are physical and approximations may result in inconsistencies. A typical example is the residual or transverse Gauß' law (2.40), which is a constraint on the physical Hilbert space. Such constraints must either be imposed on the states or one can also use them to eliminate "unphysical" degrees of freedom (here $\vec{\partial}_\perp^2 \int dx^- A^+$). The abovementioned, incomplete cancelation of IR singularities in the Tamm-Dancoff approximation occurs because the transverse Gauß' law (2.40) is violated. A more thorough discussion on this subject and possible caveats can be found in Refs. [LN+ 94a, KP 93].

Chapter 3

The Light-Front Vacuum

3.1 The Physical Picture

In the Fock space expansion one starts from the vacuum as the ground state and constructs physical hadrons by successive application of creation operators. In an interacting theory the vacuum is in general an extremely complicated state and not known *a priori*. Thus, in general, a Fock space expansion is not practical because one does not know the physical vacuum (i.e. the ground state of the Hamiltonian). In normal coordinates, particularly in the Hamiltonian formulation, this is a serious obstacle for numerical calculations. As is illustrated in Table 3.1, the LF formulation provides a dramatic simplification at this point. While all components of the momentum in normal coordinates can be positive as well as negative, the longitudinal LF momentum P_- is always positive. In free field theory (in normal coordinates as well as on the LF) the vacuum is the state which is annihilated by all annihilation operators a_k. In general, in an interacting theory, excited states (excited with respect to the free Hamiltonian) mix with the trivial vacuum (i.e. the free field theory vacuum) state resulting in a complicated physical vacuum. Of course, there are certain selection rules and only states with the same quantum numbers as the trivial vacuum can mix with this state; for example, states with the same momentum as the free vacuum ($\vec{P} = 0$ in normal coordinates, $P_- = 0$, $\vec{P}_\perp = 0$ on the LF). In normal coordinates this has no deep consequences because there are many excited states which have zero momentum. On the LF the situation is completely different. Except for pure zero-mode excitations, i.e. states where only the zero-mode (the mode with $k_- = 0$) is excited, all excited states have positive longitudinal momentum P_-. Thus only these pure zero-mode excitations can mix with the trivial LF vacuum. Thus with the exception of the zero-modes the physical LF vacuum

normal coordinates	light-front
free theory	
$P^0 = \sqrt{m^2 + \vec{P}^2}$ 	$P_+ = \frac{m^2 + \vec{P}_\perp^2}{2P_-}$ 
$P^0 = \sum_{\vec{k}} a^\dagger_{\vec{k}} a_{\vec{k}} \sqrt{m^2 + \vec{k}^2}$	$P_+ = \sum_{k_-, \vec{k}_\perp} a^\dagger_{k_-, \vec{k}_\perp} a_{k_-, \vec{k}_\perp} \frac{m^2 + \vec{k}_\perp^2}{2k_-}$
vacuum (free theory)	
$a_{\vec{k}}\|0\rangle = 0$	$a_{k_-, k_\perp}\|0\rangle = 0$
vacuum (interacting theory)	
many states with $\vec{P} = 0$ (e. g. $a^\dagger_{\vec{k}} a^\dagger_{-\vec{k}}\|0\rangle$)	$k_- \geq 0$
$\hookrightarrow \|\tilde{0}\rangle$ very complex	$\hookrightarrow$ only pure zero-mode excitations have $P_- = 0$
	$\hookrightarrow \|\tilde{0}\rangle$ can only contain zero-mode excitations

Table 3.1: Zero Modes and the Vacuum

(i.e. the ground state) of an interacting field theory must be trivial.[1]

Of course, this cannot mean that the vacuum is entirely trivial. Otherwise it seems impossible to describe many interesting problems which are related to spontaneous symmetry breaking within the LF formalism. For example one knows that chiral symmetry is spontaneously broken in QCD and that this is responsible for the relatively small mass of the pions — which play an important role in strong interaction phenomena at low energies. What it means is that one has reduced the problem of finding the LF vacuum to the problem of understanding the dynamics of these zero-modes.

First this sounds just like merely shifting the problem about the structure of the vacuum from nonzero-modes to zero-modes. However, as the free dispersion relation on the LF,

$$k_+ = \frac{m^2 + \vec{k}_\perp^2}{2k_-}, \tag{3.1}$$

indicates, zero-modes are high energy modes! Hence it should, at least in principle, be possible to eliminate these zero-modes systematically giving rise to an effective LF field theory [LT 91].

Before we embark on theoretically analyzing zero-modes, it should be emphasized that zero-modes may have experimentally measurable implications. This is discussed in Refs.[Bu 92c, Bu 95].

3.2 Examples for Zero Modes

Usually, in the context of LF quantization, fields that do not depend on x^- are called zero-modes (regardless whether they depend on $\vec{x}_\perp$ or not). However, for practical purposes, the following classification scheme seems to be particularly useful [KP 93]: If one denotes

$$\langle f \rangle_o \equiv \frac{1}{2L} \int_{-L}^{L} dx^- f(x^-, \vec{x}_\perp), \tag{3.2}$$

then

$$\langle f \rangle \equiv \frac{1}{(2L_\perp)^2} \int d^2x_\perp \langle f \rangle_o = \frac{1}{2L(2L_\perp)^2} \int d^2x_\perp \int_{-L}^{L} dx^- f(x^-, \vec{x}_\perp) \tag{3.3}$$

is called the *global zero-mode*, while

$$\overset{o}{f} \equiv \langle f \rangle_o - \langle f \rangle \tag{3.4}$$

[1] Cases where the LF Hamiltonian has no ground state will be discussed below.

is called *proper zero-mode*. The "rest", i.e.

$$\overset{n}{f} \equiv f - \langle f \rangle_0 \tag{3.5}$$

is called the *normal mode* part of f. The motivation for this distinction arises primarily from the fact that usually only the global zero-mode can develop a vacuum expectation value but also since proper and global zero-modes have a very different dynamics.

Zero modes occur in various contexts and it is not yet entirely clear to what extend the various zero-mode effects, which will be discussed below, are connected.

3.2.1 Constant Scalar Fields

In ϕ^4 theory,

$$\mathcal{L} = \frac{1}{2}\partial_\mu \phi \partial^\mu \phi - \frac{m^2}{2}\phi^2 - \frac{\lambda}{4!}\phi^4, \tag{3.6}$$

if one chooses the "wrong" sign for the mass ($m^2 < 0$, $\lambda > 0$), spontaneous symmetry breaking occurs already at the classical level. The field ϕ develops a vacuum expectation value and the symmetry $\phi \to -\phi$ is spontaneously broken. At least in $1+1$ and $2+1$ dimensions, with appropriate values for the renormalized mass, a similar behavior is observed in the quantum version. Clearly, such a scenario requires a zero-mode. In the case of ϕ^4 theory, one may imagine that a redefinition

$$\phi \to \tilde{\phi} + \langle 0|\phi|0\rangle \tag{3.7}$$

eliminates the VEV of the global zero-mode [HV 87]. However, this does not mean that one has eliminated the zero-modes. In fact, by integrating the equations of motion over x^-, one finds

$$0 = m^2\langle\phi\rangle + \frac{\lambda}{3!}\frac{1}{2L}\int_{-L}^{L} dx^- \phi^3 \ , \tag{3.8}$$

which relates the zero-mode part of the field to the normal mode part. Clearly, this nonlinear operator identity implies that (for finite L), a mere shift of the scalar field is not sufficient to completely eliminate the zero-mode. Instead, two main classes of approaches are being used get the zero-modes under control. In DLCQ one attempts to solve the zero-mode constraint equation [Eq.(3.8)] using various approximation or expansion schemes [PV 93, PV 94, PV+ 95]. Due to nonlinear effects and operator ordering ambiguities, solving Eq.(3.8) becomes a nontrivial endeavor

[PV 93, PV 94, PV+ 95, Ro 93, HK+ 92a]. In the other approach (the effective LF-Hamiltonian approach, which will be discussed in detail in Section 3.3) one makes use of the fact that zero-modes freeze out for $L \to \infty$. Instead of keeping zero-modes explicitly, one allows for an effective Hamiltonian, which should account for their effects on normal modes in the large volume limit [Bu 93].

So far it is not known whether either one of these approaches to LF-quantization (explicit zero-modes and effective LF-Hamiltonian) leads to a consistent formulation of ϕ^4 theory in the broken phase. It is also not known to what extend the particle spectrum in the equal time formulation agrees with the spectrum on the LF. Since the broken phase of ϕ^4 in 1+1 dimensions has a rather rich spectrum: mesons, solitons [2], bound states and scattering states in the soliton-antisoliton sector [3], this seems to be an ideal test case for the various approaches to scalar zero-modes on the LF. So far, all works on ϕ^4 on the LF have concentrated on demonstrating that spontaneous symmetry breaking occurs and on reproducing the numerical value of the critical coupling constant from ET quantization.[4]

One of the most striking consequences of spontaneous symmetry breaking in ϕ^4_{1+1} is the emergence of solitons. While most LF workers choose boundary conditions that make it impossible to study solitons, soliton-antisoliton scattering states are often still possible. These states often have a very clear signature [Bu 93] and one can easily determine their threshold. Considering the extensive literature on LF-ϕ^4_{1+1} (see e.g. Refs. [PV 93, PV 94, PV+ 95] and references therein), it is surprising that solitons have been ignored so far.

3.2.2 Fermionic Zero Modes

Consider the free Dirac equation

$$0 = (i\gamma_\mu \partial^\mu - M)\,\psi = \left(i\gamma_- \partial_+ + i\gamma_+ \partial_- - i\vec{\gamma}_\perp \vec{\partial}_\perp - M\right)\psi. \tag{3.9}$$

Multiplying Eq. (3.9) with $\mathcal{P}^{(+)}$, where $\mathcal{P}^{(\pm)} = \gamma^\mp \gamma^\pm/2$ are the projection matrices introduced in Section 2.3.2, one obtains

$$i\gamma^- \partial_- \psi^{(-)} = \left(i\vec{\gamma}_\perp \vec{\partial}_\perp + M\right)\psi^{(+)} \tag{3.10}$$

with $\psi^{(\pm)} = \mathcal{P}^{(\pm)}\psi$. Clearly Eq.(3.10) is a constraint equation and one must eliminate $\psi^{(-)}$ before one can canonically quantize the theory (the kinetic

[2] Often excluded by boundary conditions on the fields.

[3] In general not excluded by boundary conditions.

[4] In view of the nontrivial renormalization effects on the LF (see Chapter 4), comparing critical coupling constants on the LF with those from ET quantization is very treacherous.

term in the fermionic Lagrangian does not contain a LF-time derivative of $\psi^{(-)}$). For all modes but the zero-modes this is straightforward. However, Eq.(3.10) does not determine the x^--independent components of $\psi^{(-)}$. In other words, because of possible "integration constants", there is some ambiguity in defining the inverse of the differential operator ∂_-.

For scalar fields, the time derivative is always accompanied by a space derivative (kinetic term: $\phi\partial_+\partial_-\phi$). Therefore, the zero-mode for scalar fields is not a dynamical degree of freedom, since its time derivative does not enter the Lagrangian. For Dirac fields this is different, since there the Lagrangian is linear in the derivatives, and the fermionic zero-mode is a dynamical degree of freedom. Little is known in this case beyond perturbation theory (see e.g. Refs. [BL 91b, RM 92]).

3.2.3 Gauge Field Zero-Modes

For practical reasons one would like to work in the LF gauge $A_- = 0$ when quantizing gauge fields on the LF. The reason is that, only in the LF gauge are canonical field momenta simple. For example, in QCD, the kinetic term for the gauge field in the Lagrangian, $-\frac{1}{4}F_{\mu\nu}F^{\mu\nu}$ contains terms like $\left[D_+, \vec{A}_\perp\right]\left[D_-, \vec{A}_\perp\right]$, i.e. in general, the term multiplying $\partial_+\vec{A}_\perp$ contains interactions. As usual in LF coordinates, the canonically conjugate momentum satisfies a constraint equation

$$\vec{\Pi}_\perp = \frac{\delta\mathcal{L}}{\delta\partial_+\vec{A}_\perp} = \partial_-\vec{A}_\perp - ig\left[A_-, \vec{A}_\perp\right]. \tag{3.11}$$

Only in the LF gauge is the constraint equation for $\vec{\Pi}_\perp$ linear in the fields, and one obtains simple commutation relations between the fields.

The problem with the LF gauge, as with axial gauges in general, has to do with infrared singularities, particularly in the nonabelian case. In order to arrive at a well defined formulation of the theory, it is often very helpful to formulate the theory in a finite 'box' with periodic boundary conditions (i.e. a torus). That way, it is generally easier to keep track of surface terms that appear in formal manipulations which include integrations by parts.

If one starts from an arbitrary gauge field configuration on a torus, it is in general not possible to reach the LF gauge (or spatial axial gauges) by means of a gauge transformation [Ma 85, KP 93]. This can be easily shown by considering the Wilson loop around the torus in the x^- direction: $W = P\exp(ig\oint dx^- A_-)$. This is a gauge invariant quantity and thus does not change under a gauge transformation. If it were possible to reach the LF gauge, $A_- = 0$, be means of a gauge transformation this would mean

transforming W to 1, which is a contradiction. It turns out that on a torus, the closest one can get to the LF gauge is $\partial_- A_- = 0$, i.e. the zero-modes for A_- remain and, due to their relation to the Wilson loop around the torus, they have a gauge invariant meaning [KP 93]. They are dynamical degrees of freedom (their ∂_+ derivative enters the Lagrangian). The zero-modes of a^i behave very similar to a scalar field, in the sense that their time derivative does not enter the Lagrangian and hence they are not dynamical degrees of freedom. Recently, Kalloniatis, Robertson and collaborators [KP 93] have developed a systematic scheme to disentangle and resolve the various zero-mode problems that appear in QED and QCD. For example, projecting the QED Maxwell equations onto the proper zero-mode sector, they obtain:

$$-\partial_\perp^2 \overset{o}{A}{}^+ = g \overset{o}{J}{}^+ \tag{3.12}$$

$$-2\partial_+^2 \overset{o}{A}{}^- - \partial_\perp^2 \overset{o}{A}{}^- - 2\partial_i\partial_+ \overset{o}{A}{}^i = g\partial_\perp^2 \overset{o}{J}{}^- \tag{3.13}$$

$$-\partial_\perp^2 \overset{o}{A}{}^+ + \partial_i\partial_+ \overset{o}{A}{}^+ + \partial_i\partial_j \overset{o}{A}{}^j = g\partial_\perp^2 \overset{o}{J}{}^j \ , \tag{3.14}$$

where J^μ is the fermionic current operator. The first of these equations (3.12) is a constraint equation and can be used to eliminate the proper zero-mode of A^+ in terms of the current J^+ [5]

$$\overset{o}{A}{}^+ = -g\frac{1}{\partial_\perp^2} \overset{o}{J}{}^+ \ , \tag{3.15}$$

which again demonstrates that $A^+ = 0$ is in general not consistent with the equations of motion.

Further simplification can be obtained by taking the (transverse) divergence of Eq.(3.14), yielding

$$\partial_+ \overset{o}{A}{}^+ = g\frac{1}{\partial_\perp^2}\partial_i \overset{o}{J}{}^i \ . \tag{3.16}$$

Inserting this back into Eq.(3.14), one finds

$$-\partial_\perp^2 \left(\delta_{ij} - \frac{\partial_i\partial_j}{\partial_\perp^2}\right) \overset{o}{A}{}^j = g\left(\delta_{ij} - \frac{\partial_i\partial_j}{\partial_\perp^2}\right) \overset{o}{J}{}^j \ , \tag{3.17}$$

which can be used to eliminate the transverse projection of the proper zero-mode of A^j. Note that so far we have not yet completely fixed the gauge, since $\partial_- A^+$ sill leaves the freedom of purely transverse gauge transformations,

[5] In the charge neutral sector, the global zero-mode of J^+ vanishes and thus the inverse Laplace is well defined.

$A^\mu \to A^\mu + \partial^\mu \Omega$, where $\Omega = \Omega(x^+, x_\perp)$. One can use this residual gauge freedom to set $\partial_i \overset{o}{A}{}^i = 0$. In combination with Eq.(3.17), this completely determines the proper zero-mode of A^i. Up to this point, it seems that the zero-modes in QED pose no real problems in the LF formulation.

The real problems in this formalism arise when one tries to implement these results in a quantum formulation that includes fermions. This can be seen when one inserts the solution for $\int dx^- A_-$ back into the Lagrangian, yielding a four Fermi interaction term of the form

$$\Delta\mathcal{L} \propto \frac{1}{2L} \overset{o}{J}{}^+ \frac{1}{\partial_\perp^2} \overset{o}{J}{}^- \ . \tag{3.18}$$

Similarly, inserting the solution for $\overset{o}{A}{}^i$ into the Lagrangian yields

$$\Delta\mathcal{L} \propto \frac{1}{2L} \overset{o}{J}{}^i \frac{\delta_{ij} - \frac{\partial_i \partial_j}{\partial_\perp^2}}{\partial_\perp^2} \overset{o}{J}{}^j \ . \tag{3.19}$$

The presence of such terms, which contain the "bad" current $j_+ = \sqrt{2} e \psi^\dagger_{(-)} \psi_{(-)}$ leads to nonlinear constraint equations for $\psi_{(-)}$. Because of the difficulties in solving this nonlinear constraint equation, it has so far not been possible to write down the LF Hamiltonian for QED or QCD in terms of physical degrees of freedom and including all zero-modes, in closed form. Only perturbative expressions for the Hamiltonian in terms of physical degrees of freedom have been found so far [KP 93]. Similar problems arise in the DLCQ formulation of QCD with additional complications arising from the difficulties in quantizing the gauge field when $A^+ \neq 0$., arising from the nonlinear constraint relation between fields and their canonical momenta (3.11).

From the practitioner's point-of-view, it would be helpful to know to what extend this elaborate machinery is actually necessary if one is interested only in the large volume limit. On a finite interval, gauge field zero-modes clearly play an important role. For example, they are essential to generate the correct potential for a heavy quark-antiquark pair in 1+1 dimensions on a circle in Coulomb gauge [ES 95, KP+ 94]. However, in the latter example, zero-mode effects for color singlet states disappear in the limit of a large interval. Unfortunately, it is not clear whether this result carries through to higher dimensional gauge theories.

3.2.4 Perturbative Zero-Modes

The zero-modes discussed are either connected to purely nonperturbative effects (like in the case of spontaneous symmetry breaking for scalar fields)

or seem to be at least connected with nonperturbative physics (like infrared singular long range effects for gauge fields). There are, however, plenty of examples where zero-mode effects appear already on the level of perturbation theory. Examples include disconnected vacuum diagrams [CM 69], "generalized tadpoles" for self-interacting scalar fields [Gr 92a, Bu 93] as well as "rainbow diagrams" for the fermion self-energy [BL 91a, BL 91b]. These examples will be discussed in more detail in Chapter 4.

In perturbation theory in LF gauge the gauge field propagator

$$D_{\mu\nu}(k) = \frac{g_{\mu\nu} - \frac{k_\mu n_\nu + k_\nu n_\mu}{k\cdot n}}{k^2 + i\varepsilon} \tag{3.20}$$

$(n\cdot A = A_-)$ becomes singular as $k_- \to 0$. There exist several "prescriptions" to handle this singularity. The most useful prescription for perturbative calculations is the Mandelstam-Leibbrandt (ML) prescription [Ma 83], where one replaces

$$\frac{1}{k\cdot n} \equiv \frac{1}{k_-} \overset{\text{ML}}{\longrightarrow} \frac{1}{k_- + i\varepsilon \text{sign}(k_+)} = \frac{k_+}{k_+k_- + i\varepsilon}. \tag{3.21}$$

The crucial property of this prescription is that the pole structure is similar to the one of a typical Feynman propagator, with poles in the second and fourth quadrant of the complex k_0-plane, and thus allows to perform a Wick rotation. This is not the case for the principal value (PV) prescription

$$\frac{1}{k_-} \overset{\text{PV}}{\longrightarrow} \frac{1}{2}\left(\frac{1}{k_- + i\varepsilon} + \frac{1}{k_- - i\varepsilon}\right) \tag{3.22}$$

with poles in the first and fourth quadrant.

One of the major disadvantages of the ML prescription is the fact that it introduces additional energy (k_+) dependencies in the propagator, which cannot be generated by a canonical LF Hamiltonian [RM 94]. However, recently the ML prescription has been successfully implemented in a LF Bethe-Salpeter approach to bound states [LS 93]. Conversely, in $\text{QCD}_{1+1}(N_C \to \infty)$ the ML prescription [Wu 77] yielded a spectrum that disagreed with the canonical LF approach [Ho 74] as well as with the result from equal time quantization [Li 86]. More recently, light-like Wilson loops in 1+1 dimensions have been calculated, using various prescriptions for gauge field propagator [BD+ 94], and it was found that only the principal value prescription yields the exact area law one expects for gauge fields in 1+1 dimensions (on noncompact manifolds).

3.3 Zero Modes and the Vacuum in ε-Coordinates

3.3.1 General Considerations

For a free particle $p_+ = \frac{L}{2\varepsilon}\left(-p_- + \sqrt{p_-^2 + 2\varepsilon m^2/L}\right)$ and p_- is no longer restricted to positive values (Fig. 2.1). Therefore, for all finite values of ε/L, the vacuum in ε-coordinates is nontrivial. Since ε-coordinates (see Section 2.2) provide a controlled and well defined approach to the LF, it seems very natural to employ this framework for studying the LF vacuum.

Let us first consider the canonical LF limit (L fixed, $\varepsilon \to 0$). In this case it is straightforward to derive an effective LF-Hamiltonian from the ε-Hamiltonian [LT 91] (for a related work, see Refs. [PN+ 95, VF 94]). For finite L the momenta are discrete. Without interactions the energy of the zero-mode ($p_+(0) = m\sqrt{\frac{L}{2\varepsilon}}$) and the energy of modes with negative momenta ($p_+(-n) \approx \frac{2\pi n}{\varepsilon}$) diverge as $\varepsilon \to 0$, while the energy of all positive momentum modes ($p_-(n) \approx \frac{m^2 L}{4\pi n}$) remains finite. For interacting fields there will be some slight quantitative changes, but the general picture should remain the same: zero-modes and negative momentum modes are high energy modes — separated from positive momentum modes by an energy gap of $\mathcal{O}\left(\sqrt{\frac{1}{\varepsilon}}\right)$ and $\mathcal{O}\left(\frac{1}{\varepsilon}\right)$ respectively. Thus although $p_- \leq 0$ modes may acquire nontrivial occupations, $p_- > 0$ modes have too little energy to cause any excitations within the $p_- \leq 0$ sector for $\varepsilon \to 0$: the $p_- \leq 0$ modes freeze out and can be replaced by their vacuum expectation value (VEV).

At this point it seems that we have succeeded in deriving a nontrivial effective LF-Hamiltonian. Unfortunately, we arrived at this result by approaching the LF in such a way that the invariant length of the interval ($\propto L\varepsilon$) approaches zero, i.e. as discussed in Ref. [LT 91], the effective theory that we have obtained is not necessarily equivalent to the original covariant theory. This can be easily illustrated by means of a perturbative example. Consider a simple tadpole with a mass insertion (to make it convergent) in $1+1$ dimensions

$$\tilde{\Sigma} = \int \frac{d^2k}{(2\pi)^2} \frac{1}{\left(k^2 - m^2 + i0\right)^2} = \frac{i}{4\pi m^2}. \tag{3.23}$$

On a finite interval (with ε coordinates) one obtains instead ($k_-(n) = \frac{2\pi}{L}n$)

$$\tilde{\Sigma} \;=\; \frac{1}{L}\sum_{k_-} \int \frac{dk_+}{2\pi} \frac{1}{\left(\frac{2\varepsilon}{L}k_+^2 + 2k_+k_- - m^2 + i0\right)^2}$$

$$= \frac{i}{4\sqrt{2\varepsilon L}} \sum_{n=-\infty}^{\infty} \left[\frac{(2\pi n)^2}{2\varepsilon L} + m^2 \right]^{-3/2} . \tag{3.24}$$

Clearly, in order to recover the continuum result (3.23) one must take limits in such a way that the invariant length if the interval becomes infinite. If one takes the LF limit first ($\varepsilon \to 0$, L fixed), one obtains a divergent contribution from the zero-mode.

A different result is obtained if one performs the continuum limit first ($L \to \infty$, ε/L fixed). Since this corresponds to $\varepsilon L \to \infty$ no problems with perturbation theory arises. However, since the spectrum is now continuous, there is no mass gap and the derivation of the effective Hamiltonian for $\varepsilon/L \to 0$ becomes more complicated. Nevertheless, it is still possible: first, note that the momentum scale of the continuum Hamiltonian is $m^2\varepsilon/L$ since momentum dependent terms in the continuum Hamiltonian appear only in the kinetic term $\propto \frac{L}{2\varepsilon}\left[-k_- + \sqrt{k_-^2 + m^2 2\varepsilon/L}\right]$ and in vertex factors $\propto \left(k_-^2 + m^2 2\varepsilon/L\right)$. Thus the typical momentum scale in the vacuum is given by $p_-^{vac} = \mathcal{O}\left(\sqrt{\frac{\varepsilon}{L}} m\right)$. Similarly the energy scale for vacuum excitations (zero total P_-) is of the order $\mathcal{O}\left(\sqrt{\frac{L}{\varepsilon}} m\right)$. Suppose one is interested in the effective Hamiltonian for a physical particle of total momentum p_-^{tot} moving in the vacuum. If $\frac{\varepsilon}{L} \ll 1$ then there is almost no overlap between the wave function of the vacuum $p_-^{vac} = \mathcal{O}\left(\sqrt{\frac{\varepsilon}{L}} m\right)$ and the wave function of the partons in the particle p_-^{parton} because the parton wavefunction (calculated for example with a typical LF Hamiltonian) vanishes for small momenta. Thus one can introduce an energy gap *by hand* without affecting the dynamics in the limit $\frac{\varepsilon}{L} \to 0$: for example, by selecting cutoffs Λ_1 and Λ_2 such that

$$m\sqrt{\frac{\varepsilon}{L}} \ll \Lambda_1 \ll \Lambda_2 \ll p_-^{tot} \tag{3.25}$$

and removing all modes with $\Lambda_1 < k_- < \Lambda_2$. First, this gives rise to a mass gap and one can argue that the modes with $k_- < \Lambda_1$ remain frozen (energy scale $k_+ > \frac{m^2}{2\Lambda_1}$) when excitations with $k_- > \Lambda_2$ are present (energy scale $k_+ < \frac{m^2}{2\Lambda_2}$): in second order perturbation theory, the energy shift for modes with $k_- > \Lambda_2$ due to excitations of n modes with $k_- < \Lambda_1$ is given by

$$\Delta^{(2)} E \propto \frac{\left|\Lambda_1^{-n/2}\right|^2}{-\frac{1}{\Lambda_1}} \Lambda_1^n = -\Lambda_1 \overset{\Lambda_1 \to 0}{\longrightarrow} 0 \tag{3.26}$$

Here $\Lambda_1^{-n/2}$ is a vertex factor, arising from the factor $\frac{1}{\sqrt{\omega_n}}$ in the expansion of the fields ϕ (Eq.(2.10)), the factor Λ_1^n is the phase space factor for n modes

with $k_- = \mathcal{O}(\Lambda_1)$ and states with $k_- < \Lambda_1$ excitations are off-shell by at least $\frac{1}{\Lambda_1}$.

Since $k_- < \Lambda_1$ excitations are suppressed, the effective LF-Hamiltonian for the modes with $k_- > \Lambda_2$ contains the $k_- < \Lambda_1$ modes only via their VEV (which may be nontrivial!)

$$V^{eff}_{k_->\Lambda_2} = \langle 0_{k_-<\Lambda_1} | V | 0_{k_-<\Lambda_1} \rangle. \tag{3.27}$$

The crucial point is that the parton distribution calculated with such an effective LF Hamiltonian vanishes for small momenta in the above super-renormalizable example [6]. Thus as long as Λ_2 is small enough compared to the total momentum of the particle p, the parton distribution vanishes already for momenta much larger than Λ_2 and the presence of the cutoff does not affect the parton dynamics. Since the VEVs are nearly independent of $\frac{\varepsilon}{L}$, so is the effective Hamiltonian. Thus the suppression of the parton distribution due to the kinetic energy sets in at a value cp^{tot}_-, where c is nearly independent of $\frac{\varepsilon}{L}$. Thus, for $\frac{\varepsilon}{L} \to 0$, Λ_2 can easily be chosen smaller than cp^{tot}_- while Eq.(3.25) remains satisfied. In other words, Λ_2 can be chosen such that the parton dynamics is independent of the exact position of the cutoff. Similarly, since vacuum momenta are restricted to $p^{vac}_- = \mathcal{O}\left(\sqrt{\frac{\varepsilon}{L}} m\right) \ll \Lambda_1$, the presence of the cutoff does not affect the dynamics of the vacuum either, i.e., the numerical value of the VEVs which enter Eq.(3.27) is independent of the cutoff.

In the 2nd order perturbation theory argument above we made use of $\Lambda_1 \ll \Lambda_2$ to make sure that the energy denominator in Eq.(3.26) is of the order $\mathcal{O}\left(\frac{1}{\Lambda_1}\right)$. This is actually not necessary, since the occupation of these modes is anyway *dynamically* suppressed for $k_- < cp_-$ and as long as $\Lambda_1 \ll cp_-$, the energy denominator will automatically be $\mathcal{O}\left(\frac{1}{\Lambda_1}\right)$ or smaller.

Thus we can actually let $\Lambda_2 \to \Lambda_1$, i.e. remove the cutoff, without altering the conclusion. Introducing a mass gap was helpful in deriving an effective Hamiltonian for modes with $p_- \gg m\sqrt{\frac{\varepsilon}{L}}$. However, since the solutions of the effective Hamiltonian vanish at small p_- anyway, there is no need for a cutoff: a region void of excitations between $m\sqrt{\frac{\varepsilon}{L}}$ and cp_- develops dynamically (Figure 3.1) and this is sufficient to derive an effective Hamiltonian.

In the end, the following result is obtained. Suppose we started from some polynomial interaction

$$\mathcal{L}^{int} = -\sum_n \lambda_n \frac{\phi^n}{n!}. \tag{3.28}$$

[6] Roughly speaking, the LF kinetic energy T, which one can calculate from the parton momentum distribution $f(k_-)$, using $T = m^2 \int_0^{P^{tot}_-} dk_- \frac{f(k_-)}{2k_-}$, has to remain finite.

Figure 3.1: Schematic occupation of modes in the presence of a particle with momentum p_- for $\varepsilon/L \ll 1$ (i.e. "close to the LF"). The modes near $k_- = 0$ are already present in the vacuum and are dynamically restricted to $k_- = \mathcal{O}\left(m(\varepsilon/L)^{(1/2)}\right)$ The "parton distribution", i.e. the modes which are occupied in the presence of the particle but not in the vacuum, vanish at small k_- at a momentum scale which remains finite as $\varepsilon/L \to 0$. The presence of the cutoffs has almost no effect on the dynamics.

Then, using Eq.(3.27) (after some combinatorics) the effective interaction, which enters the LF Hamiltonian for $p_- > 0$ modes in the limit $\frac{\varepsilon}{L} \to 0$ is given by

$$\mathcal{L}_{eff}^{int} = -\sum_n \lambda_n \sum_{k=0}^{n} \frac{\phi^{(n-k)}}{(n-k)!} \langle 0| \frac{\phi^k}{k!} |0\rangle \tag{3.29}$$

(in order to obtain this result one also uses that, after normal ordering, the $p_- > 0$ modes do not contribute to the VEVs). Eq.(3.29) is a remarkable result. It states that nontrivial vacuum effects enter the LF-Hamiltonian only via effective interactions. The effective coupling constants depend on the vacuum condensates which, in general, cannot be obtained directly from a LF calculation [7]. They must be considered as renormalization parameters of the LF theory. Eq.(3.29) is also valid in situations where spontaneous symmetry breaking occurs. For example in ϕ^4 theory, $\langle 0|\phi|0\rangle$ may become nonzero and a ϕ^3 interaction will thus appear in the effective Lagrangian. [8] However, note that only a *finite* number of condensates is necessary to specify the effective LF Hamiltonian: if N is the highest power of ϕ entering

[7] However, there are exceptions where one can use sum rules or consistency conditions to determine the effective couplings iteratively. Examples will be discussed in the following section.

[8] It should be emphasized that we did not make any mean field assumptions, such as $\langle 0|\phi^k|0\rangle = \langle 0|\phi|0\rangle^k$, in order to arrive at this result.

the canonical LF Hamiltonian then only condensates $\langle 0|\phi^k|0\rangle$ with $k < N$ need to be considered.

At several points in the above discussion it was important that the theory is free of divergences (up to a finite number of diagrams which can always be subtracted before applying above argumentation). First this was important to insure that the momentum scale in the vacuum is finite. Secondly, it was important because only in the absence of divergences one can apply the kinetic energy argument to prove that the parton momentum distribution vanishes for small parton momenta. Therefore one must be very careful when generalizing the above results to higher dimensional field theories. Eq.(3.29) can be used in renormalizable field theories only *after* a cutoff has been imposed. This is for example the case for the transverse lattice which will be discussed in Section 5.3.

3.3.2 A simple Example for the limit $L \to \infty$, $\varepsilon/L \to 0$

The appearance of the gap as $L \to \infty$ (first) and $\varepsilon/L \to 0$ is best understood by studying a concrete example. Ideally this implies considering some non-trivial interacting field theory and calculating the occupation of the modes nonperturbatively for various ε/L and $L \to \infty$. However, even in integrable models, such as the sine-Gordon model, the occupation of the modes is not known exactly! Since numerical calculations at small but finite ε/L and $L \to \infty$ are very complicated — particularly if one is interested in momenta of the order of $m\sqrt{\varepsilon/L}$ — and we will proceed by studying a perturbative example. Due to the fact that the appearance of the gap is mostly a consequence of dimensional analysis, this will be sufficient to highlight the essential physics of the limit $\varepsilon/L \to 0$, $L \to \infty$ ($L \to \infty$ first). The example which we will consider is a scalar field theory in $1+1$ dimensions with polynomial self-interactions

$$\mathcal{L} = \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - \frac{m^2}{2}\phi^2 - \frac{\lambda_4}{4!}\phi^4 - \frac{\lambda_6}{6!}\phi^6. \tag{3.30}$$

In ε-coordinates the Hamiltonian for this model reads after normal ordering

$$H = \sum_n p_+(k_n) a^\dagger_{k_n} a_{k_n} + \int_0^L dx^- \left[\frac{\lambda_4}{4!} : \phi^4 : + \frac{\lambda_6}{6!} : \phi^6 :\right], \tag{3.31}$$

where the same notation as in Section 2.2 has been used. In lowest (zeroth) order in λ_4 and λ_6 the vacuum is the Fock vacuum, defined by $a_{k_n}|0\rangle = 0$. This changes of course for nonvanishing couplings. For example, in second order perturbation theory in λ_4 one finds for the occupation of states in the

vacuum

$$\begin{aligned}\rho_0(k,\frac{\varepsilon}{L},L) &\equiv \langle\tilde{0}|a_k^\dagger a_k|\tilde{0}\rangle \\ &= \frac{\lambda_4^2}{3!}L^2\sum_{k_2,\ k_3}\frac{1}{\left[p_+(k)+p_+(k_2)+p_+(k_3)+p_+(k_4)\right]^2} \\ &\quad\times\frac{1}{2\omega(k)}\frac{1}{2\omega(k_2)}\frac{1}{2\omega(k_3)}\frac{1}{2\omega(k_4)},\end{aligned} \tag{3.32}$$

where $\omega(q) = L\sqrt{q^2+2\varepsilon m^2/L}$, $p_+(q) = \left(-q+\sqrt{q^2+2\varepsilon m^2/L}\right)L/2\varepsilon$ and $k_4 = -k-k_2-k_3$. A similar expression is found for the $\mathcal{O}(\lambda_6^2)$-term, which will be omitted for simplicity. In the limit $L\to\infty$ one thus finds

$$\begin{aligned}\rho_0(k,\frac{\varepsilon}{L}) &\equiv \lim_{L\to\infty}\frac{L}{2\pi}\frac{1}{\sqrt{2\varepsilon L m^2}}\rho_0(k,\frac{\varepsilon}{L},L) \\ &= \frac{\lambda_4^2}{96\pi m^5}\sqrt{\frac{L}{2\varepsilon}}\hat{\rho}_0\left(k\sqrt{\frac{L}{2\varepsilon m^2}}\right),\end{aligned} \tag{3.33}$$

where

$$\hat{\rho}_0(z) = \int_{-\infty}^{\infty}\frac{dz_2}{2\pi}\int_{-\infty}^{\infty}\frac{dz_3}{2\pi}\frac{\hat{\omega}(z)^{-1}\hat{\omega}(z_2)^{-1}\hat{\omega}(z_3)^{-1}\hat{\omega}(z_4)^{-1}}{\left[\hat{\omega}(z)+\hat{\omega}(z_2)+\hat{\omega}(z_3)+\hat{\omega}(z_4)\right]^2} \tag{3.34}$$

with $\hat{\omega}(z) = \sqrt{z^2+1}$ and $z_4 = -z-z_2-z_3$. The factor $L/2\pi$ arises from going from discrete to continuous momentum k and we divided by the invariant length of the interval because the occupation in the vacuum trivially scales like the invariant length.

Most importantly, the momentum scale in the occupation of the vacuum is set by $m\sqrt{2\varepsilon}/L$. The momentum density in the vacuum is sharply peaked around $k=0$ with width $\mathcal{O}(\sqrt{2\varepsilon/L})$ and height $\mathcal{O}(\sqrt{L/2\varepsilon})$, i.e. it resembles a δ-function as $\varepsilon/L\to 0$.

Let us now consider a state with momentum P, where P is taken independent of L or ε. To lowest order

$$|P\rangle = a_P^\dagger|0\rangle \tag{3.35}$$

and thus

$$\rho_P(k,\frac{\varepsilon}{L},L) \equiv \langle P|a_k^\dagger a_k|P\rangle = \delta_{k,P}+\mathcal{O}(\lambda^2). \tag{3.36}$$

Three classes of corrections contribute to ρ_P: insertions in disconnected vacuum diagrams (Fig.3.2a) [yielding again Eq.(3.33)], insertions in tadpoles (Fig.3.2b) and the rest, i.e. insertions in non-tadpole connected corrections (Fig.3.2c).

Figure 3.2: $\mathcal{O}(\lambda^2)$-corrections to the mode density in the presence of a particle with momentum P. a.) disconnected corrections, b.) insertions into generalized tadpoles (i.e. diagrams where a subgraph is connected with the rest of the diagram at one point only) and c.) non-tadpole connected corrections.

The tadpole term yields

$$\begin{aligned}\tilde{\rho}_P^{tadpole}(k,\sqrt{\frac{\varepsilon}{L}}) &\equiv \lim_{L\to\infty}\frac{L}{2\pi}\rho_P^{tadpole}(k,\sqrt{\frac{\varepsilon}{L}},L)\\ &= \frac{\lambda_4\lambda_6}{96\pi m^4 P}\hat{\rho}_0\left(k\sqrt{\frac{L}{2\varepsilon m^2}}\right),\end{aligned} \tag{3.37}$$

and for the non-tadpole, connected term one finds

$$\begin{aligned}\tilde{\rho}_P^{nt}(k,\sqrt{\frac{\varepsilon}{L}}) &\equiv \lim_{L\to\infty}\frac{L}{2\pi}\rho_P^{nt}(k,\sqrt{\frac{\varepsilon}{L}},L)\\ &= \frac{\lambda_4^2}{32\pi}\int_{-\infty}^{\infty}\frac{dk_2}{2\pi}\left[\frac{1}{E_A^2}+\frac{1}{E_B^2}\right]\frac{1}{\omega(P)}\frac{1}{\omega(k)}\frac{1}{\omega(k_2)}\frac{1}{\omega(k_3)}\end{aligned} \tag{3.38}$$

($k_3 = P-k-k_2$) plus a similar term proportional to λ^6, which will be omitted in the following for simplicity. The energy denominators in Eq.(3.38), corresponding to the two time orderings, are given by

$$\begin{aligned}E_A &= p_+(P)-p_+(k)-p_+(k_2)-p_+(k_3)\\ E_B &= -p_+(P)-p_+(k)-p_+(k_2)-p_+(k_3).\end{aligned} \tag{3.39}$$

The various contributions to the occupations in the presence of a particle with momentum P are shown in Fig. 3.3 for a number of values for ε/L. The numerical values for m and P, as well as the coupling constants λ_4 and λ_6, in the plots are taken to be 1. Several effects can be observed:

Figure 3.3: $\mathcal{O}(\lambda^2)$ contributions to the occupation density $\rho(\tilde{k})$ in the presence of an excitation with momentum $P = 1$ and mass $m = 1$ for various values of the parameter ε/L. Dashed line: disconnected vacuum contribution, dotted line: tadpole contribution, full line: non-tadpole connected (dispersive) contribution.

Figure 3.4: Same as in the previous Figure but for smaller values of ε/L and plotted over a logarithmic momentum scale.

- The non-tadpole connected (dispersive) contribution scales in the limit $L \to \infty$, $\varepsilon/L \to 0$. The scaling function is the LF momentum distribution.

- Both, the disconnected contribution as well as the tadpole contribution, are restricted to a region $k^2 = \mathcal{O}(\varepsilon/L)\Lambda^2$ near the origin, where Λ is some mass scale ($\Lambda = m$ to lowest nontrivial order).

- The integral over the disconnected vacuum contribution is independent of ε/L.

- Compared to the vacuum contribution, the tadpole term is suppressed by one power of $\sqrt{\varepsilon/L}$ and thus can be neglected as $\varepsilon/L \to 0$

The gap can be most easily observed by plotting the density over a logarithmic momentum scale (Fig.3.4).

For very small values of ε/L, the momentum distributions from the disconnected diagrams [momentum scale $\mathcal{O}(\sqrt{\varepsilon/L})$] and from the dispersive contributions [momentum scale $0.1 - 1$] no longer overlap and a gap arises. The disconnected contributions were already present in the vacuum (ground state for $P = 0$) and are unaltered by the presence of the excitation with momentum P. The only change in occupation within the small momentum region arises from the tadpoles but its integrated contribution vanishes as $\varepsilon/L \to 0$ and becomes negligible in that limit. Note that $\sqrt{\varepsilon/L}$ must be extremely small for the gap to be clearly visible. This makes nonperturbative studies of the gap forbiddingly difficult numerically because one would have to cover a huge number of scales (from $\sqrt{\varepsilon/L}$ to 1) while keeping the invariant volume large.

3.4 Vacuum Condensates and Sum Rules

In the previous section we explained that vacuum condensates may enter the effective (zero-mode free) LF Hamiltonian via induced coupling constants. The condensates cannot be calculated directly unless one includes dynamical zero-modes. However, even without zero-modes, it is possible to calculate at least some of the condensates indirectly using sum rule techniques. As an example, let us consider the two point function in a self-interacting scalar field theory

$$G(x) \equiv \langle 0|\phi(0)\phi(x)|0\rangle \tag{3.40}$$

($x^0 < 0$, $x^2 < 0$). Inserting a complete set of states one obtains

$$\begin{aligned} G(x) &= \langle 0|\phi|0\rangle^2 + \sum_n \int_0^\infty \frac{dp_-}{2p_-} \langle 0|\phi(0)|n,p\rangle\langle n,p|\phi(x)|0\rangle \\ &= \langle 0|\phi|0\rangle^2 + \sum_n \int_0^\infty \frac{dp_-}{2p_-} \left|\langle 0|\phi(0)|n,p\rangle\right|^2 \exp\left(i(p_- x^- + p_+^n x^+)\right), \end{aligned} \tag{3.41}$$

where the sum is over all particle states. The normalization of the states is $\langle n,p|m,p'\rangle = 2p_-\delta(p_- - p'_-)\delta_{nm}$ and the energies are given by the on-shell dispersion relation $p_+^n = M_n^2/2p_-$. By boost invariance (in the continuum limit), the vacuum to "hadron" matrix elements are independent of the momentum

$$\langle 0|\phi|n,p\rangle = \frac{g_n}{\sqrt{2\pi}}, \tag{3.42}$$

and thus

$$G(x) = \langle 0|\phi|0\rangle^2 - \sum_n \frac{|g_n|^2}{4\pi} K_0(M_n\sqrt{-x^2}), \tag{3.43}$$

where K_0 is a modified Bessel function [AS 70]. In the limit $x^2 \to 0$ one thus finds

$$\begin{aligned} \langle 0|\phi^2|0\rangle - \langle 0|\phi^2|0\rangle_{free} &\equiv lim_{x^2\to 0}\left[G(x) - G(x)_{free}\right] \\ &= \langle 0|\phi|0\rangle^2 + \sum_n \frac{|g_n|^2}{4\pi} \log\frac{M_n}{M_{free}}, \end{aligned} \tag{3.44}$$

where we used $\sum_n |g_n|^2 = 1$ and M_{free}, $G_{free}(x)$ are the invariant mass and the two point function for noninteracting fields. Eq. (3.44) is very interesting because it allows to calculate $\langle 0|\phi^2|0\rangle$ in terms of $\langle 0|\phi|0\rangle$ and quantities (M_n and g_n) which are calculable in a canonical LF calculation without any dynamical zero-modes.

A similar trick works for the cubic condensates. Of course one has to be careful to separate the disconnected contributions first

$$\begin{aligned} \langle 0|\phi(x)\phi(y)\phi(z)|0\rangle &= \langle 0|\phi|0\rangle^3 + \langle 0|\phi(x)\phi(y)\phi(z)|0\rangle_C^3 \\ &+ \langle 0|\phi|0\rangle\left[\langle 0|\phi(x)\phi(y)|0\rangle_C + \langle 0|\phi(x)\phi(z)|0\rangle_C\right. \\ &\qquad \left. + \langle 0|\phi(y)\phi(z)|0\rangle_C\right]. \end{aligned} \tag{3.45}$$

The connected piece is calculated similar to $\langle 0|\phi^2|0\rangle$ by inserting a complete set of states. In the limit $(x-y)^2 \to 0$, $(y-z)^2 \to 0$ one finds

$$\begin{aligned} \langle 0|\phi^3|0\rangle &= \langle 0|\phi|0\rangle^3 + 3\langle 0|\phi|0\rangle\langle 0|\phi^2|0\rangle_C \\ &\quad + \sum_n \int_0^\infty \frac{dp_-}{2p_-} \langle 0|\phi|n,p_-\rangle\langle n,p_-|\phi^2|0\rangle_C \\ &= \langle 0|\phi|0\rangle^3 + 3\langle 0|\phi|0\rangle\langle 0|\phi^2|0\rangle_C \\ &\quad + \sum_n \frac{g_n h_n}{4\pi} \log M_n, \end{aligned} \tag{3.46}$$

where $h_n \equiv \sqrt{2\pi}\langle 0|\phi^2|n, p_-\rangle_C$ (independent of p_-) and $\langle 0|\phi^2|0\rangle_C$ can be taken from above (3.44). Note that $\sum_n g_n h_n = 0$ because the states $\int dp_- \exp(ip_-x^-)\phi(x^-)|0\rangle$ and $\int dp_- \exp(ip_-x^-)\phi^2(x^-)|0\rangle$ are orthogonal. Like g_n, h_n can be calculated in a LF calculation without dynamical zero-modes.

The generalization of these results to higher condensates is straightforward and by recursion one can express them in terms of $\langle 0|\phi|0\rangle$ and matrix elements which are accessible in a LF calculation. These matrix elements (g_n and h_n) depend on the states and thus implicitly on the coupling constants in the effective LF Hamiltonian. Since the coupling constants in the effective LF Hamiltonian also involve the condensates (3.29), this implies that it may be possible to determine the coupling constants in the effective LF Hamiltonian self consistently.

Similar results may be derived for Yukawa theories. In Section 4.2 we will relate the effective coupling constants in the LF Hamiltonian to the spectral densities (4.16) which are also accessible in a LF calculation.

Extracting vacuum condensates from a canonical LF calculation via sum rules has for example been done in Ref. [Zh 85] for the $m_q \to 0$ quark condensate in $\mathrm{QCD}_2(N_C \to \infty)$. The numerical result for $\bar{\psi}\psi$ was confirmed later in Ref. [Li 86] in an equal time framework. A finite quark mass calculation, based on LF wavefunctions and sum rule techniques, was first done in Refs. [Bu 89a, Bu 94b]. Again the result agreed with the result from equal time quantization [LT 89].

Chapter 4

Perturbative Renormalization

In practical applications of LF quantization, such as calculating parton distributions, nonperturbative effects play a major role. Nevertheless it makes sense to study renormalization of LF field theories first from a perturbative point of view because this allows to resolve some issues which would also appear in a nonperturbative bound state equation.

Most terms in the perturbation series generated by the LF Hamiltonian of QED or QCD are UV-divergent. This is not very surprising. After all we have become used to the fact that most quantum field theories contain divergences. However, as we will see in the following, the structure of the divergences in light front perturbation theory (LFPTh) is different from the divergences in covariant perturbation theory (CPTh). Because LF quantization is a noncovariant formulation of field theory, different Lorentz components of a divergent expression are not necessarily related to each other. In addition, in many examples the degree of divergence in LFPTh is worse than in CPTh.

On the one hand this is caused by the choice of regulators. On a formal level, LFPTh and CPTh are equivalent [CY 72]. However, the "equivalence proof" involves steps which are ill defined in the presence of divergences and singularities. In practice, if one wants to demonstrate the equivalence between LFPTh and CPTh, it is very helpful to completely regularize the theory at the level of the Lagrangian — before quantizing. One possibility to do this is Pauli-Villars regularization, where one can introduce as many regulators as are necessary to render the theory free of divergences and light-cone singularities [CY 72, BF+ 72]. Obviously, it is then not difficult to establish the equivalence between LFPTh and CPTh. However, for practical applications, Pauli-Villars regularization is not very useful. On the one hand the Hamiltonian for a Pauli-Villars regularized theory is either nonhermitian or unbounded from below or both. [1] On the other hand, Pauli-Villars regulators

[1] The Pauli-Villars ghosts, must be quantized with the "wrong" commutation relations

are not very useful for nonabelian gauge theories, because there one would have to introduce massive vector fields, which will in general destroy the renormalizability of nonabelian gauge theories. For these reasons, one is not interested in employing these regulators in the context of LF quantization.

For practical applications, it is very useful to use regulators that are compatible with the kinematic symmetries [2] of the LF.[3] In the literature one finds for example the Brodsky-Lepage regulator

$$\sum_i \frac{\vec{k}_{\perp i}^2 + m_i^2}{x_i} < \Lambda_{BL}^2, \tag{4.1}$$

where the sum extends over all particles and $x_i = k_{i-}/P_-^{tot} \in (0;1)$ are LF momentum fractions. Other regulators are a transverse momentum cutoff

$$\vec{k}_\perp^2 < \Lambda_\perp^2 \tag{4.2}$$

or dimensional regularization in the transverse direction [MP+ 91, BL 91a]

$$\int d^2k_\perp \to \int d^{2(1-\epsilon)}k_\perp. \tag{4.3}$$

Very often it is in addition necessary to introduce a cutoff for small longitudinal momenta, such as

$$\Theta(x_i - \delta) \tag{4.4}$$

and/or a cutoff in the number of particles (Tamm-Dancoff approximation). What all these regulators have in common is that they are in general not compatible with Lorentz transformations that are not kinematic symmetries of the LF (like rotations around any axis other than the z-axis). Thus when using one of these regulators, one should not be surprised if matrix elements do not exhibit the full Lorentz invariance — unless one compensates for this effect by means of a more general counterterm structure. This last point will be the main subject for the rest of this chapter. The Tamm-Dancoff approximation will be discussed in more detail in Section 5.4.

4.1 Scalar Fields

The following observation is very helpful in analyzing the perturbative equivalence between CPTh and LFPTh: Hamiltonian (with x^0 or x^+ as time) perturbation theory can be obtained from covariant perturbation theory after

in order to contribute with opposite signs in loops, which is necessary to cancel the divergences. The properties of the Hamiltonian then follow from the spin statistics theorem.

[2] These are all Poincaré transformations, which leave the $x^+ = 0$ initial surface invariant, such as translations, rotations around the $z - axis$ or longitudinal boosts.

[3] This excludes, e.g. Euclidean lattices.

Figure 4.1: 1-loop self-energy diagram in ϕ^3_{1+1}.

integrating the energies (i.e. the momentum variable which is canonically conjugate to the "time") first. Thus from the mathematical point of view, the question about equivalence between LFPTh and CPTh has been reduced to the question whether the order of integration plays a role in a Feynman integral.

As an example, let us consider the 1-loop self-energy Σ in ϕ^3-theory in 1+1 dimensions (Figure 4.1)

$$\Sigma = \frac{ig^2}{2}\int \frac{dk_- dk_+}{(2\pi)^2}\frac{1}{k^2-m^2+i\varepsilon}\frac{1}{(p-k)^2-m^2+i\varepsilon} \tag{4.5}$$

$$= \frac{g^2}{2}\int \frac{dk_-}{2\pi}\frac{\Theta(k_-)}{2k_-}\frac{\Theta(p_- - k_-)}{2(p_- - k_-)}\frac{1}{p_+ - \frac{m^2}{2k_-} - \frac{m^2}{2(p_- - k_-)}}. \tag{4.6}$$

First, without going into the details, it is easy to convince oneself that Eq.(4.6) is exactly what one obtains in LF-Hamiltonian perturbation theory: $p_+ - \frac{m^2}{2k_-} - \frac{m^2}{2(p_- - k_-)}$ is the energy denominator and the Θ-functions ensure that all momenta are positive. The other factors arise from a vertex factor proportional to $(k_-\,(p_- - k_-))^{-1/2}$ at each vertex. It is also easy to see that Eq.(4.6) agrees with the covariant calculation with symmetric integration. After substituting $k_- = xp_-$ in Eq.(4.6) one finds

$$\Sigma = \frac{g^2}{2}\int_0^1 \frac{dx}{4\pi}\frac{1}{p^2x(1-x)-\lambda^2}. \tag{4.7}$$

In the covariant calculation one first combines the two denominators in Eq.(4.5) with a Feynman parameter integral and then one integrates symmetrically over d^2k. This reproduces Eq.(4.7) where the x-integration corresponds to the parameter integral.

Our next example will be one where the order of integration does matter, namely the so called simple tadpole diagram in ϕ^4 (for simplicity again in

Figure 4.2: Typical generalized tadpole diagrams for ϕ^4

1+1 dimensions)

$$\Sigma = \frac{ig}{2} \int \frac{d^2k}{(2\pi)^2} \left(\frac{1}{k^2 - m^2 + i\varepsilon} - \frac{1}{k^2 - \Lambda^2 + i\varepsilon} \right). \tag{4.8}$$

We have already performed a subtraction because the unregularized integral diverges logarithmically. Symmetric integration over d^2k yields $\Sigma = (g/8\pi) \log \Lambda^2/m^2$. In LFPTh (unsymmetric integration; k_+-integral first) one obtains zero: for $k_- \neq 0$ one can always close a contour integral in the complex k_+ plane such that no poles are enclosed. The surface term vanishes because of the subtraction term. The point $k_- = 0$ is usually omitted in LF quantization without zero-modes. The mathematical reason for the difference between the LFPTh result and the CPTh result is a term $\propto \delta(k_-)$, which is omitted if one (as is usually, either explicitly or implicitly done) has a small k_- cutoff, like $\Theta(k_- - \varepsilon)$ at each line — even in the limit $\varepsilon \to 0$ [CM 69].

This result is very typical for pathologies of LFPTh with scalar fields. Compared to CPTh, one omits certain diagrams which are nonzero in CPTh, i.e. LFPTh yields *a priori* wrong results! Fortunately (later we will see that there is a good reason for this) the 'mistake' does not depend on the external momenta. Thus one can make up for the mistake by means of a local counterterm in the Lagrangian. Other diagrams which suffer from the same problem are the generalized tadpole diagrams, i.e. diagrams where part of the diagram is connected with the rest of the diagram only at one single point (examples are shown in Figs.4.2 and 4.3). As discussed in Ref.[Bu 93], they are all zero in LFPTh. However, because the generalized tadpoles in these diagrams are connected to the rest of the diagram only at one point, the covariant calculation yields a momentum independent result for the tadpole part (just a number), which can thus always be replaced by a local insertion

Figure 4.3: Typical tadpole diagrams arising for scalar fields with more general polynomial self-interactions

into the diagram. In practice this means that the fact that all generalized tadpoles are (wrongfully) zero on the LF, can be easily compensated by appropriate redefinitions of bare coupling constants! Furthermore, tadpole diagrams are the only diagrams which are treated incorrectly in naive LFPTh.

A very interesting result is the relation between the tadpole counterterms and vacuum condensates [Bu 93]. For example, each tadpole correction to the propagator in Fig.4.2a can be written as a mass insertion times the free field vacuum expectation value (VEV) of $\langle 0|\phi^2|0\rangle$. The generalized tadpole in Fig.4.2b corresponds to a mass insertion times a higher order correction $\langle 0|\phi^2|0\rangle$. The higher order tadpoles in Fig.4.3 correspond to mass (a) and vertex (b) insertions times a term that contributes to $\langle 0|\phi^4|0\rangle$. Suppose the interaction term in the original Lagrangian is

$$\mathcal{L}_{int} = -\sum_n \frac{\lambda_n}{n!}\phi^n. \tag{4.9}$$

Then all the 'missing tadpoles' are automatically taken into account if one uses

$$\mathcal{L}_{int,eff} = -\sum_n \lambda_n \sum_{k=0}^{n} \frac{\phi^{n-k}}{(n-k)!}\frac{\langle 0|\phi^k|0\rangle}{k!}. \tag{4.10}$$

In other words, $\mathcal{L}_{int,eff}$ with naive LFPTh yields the same results as $\mathcal{L}_{int}$ (4.9) (4.10) with CPTh to all orders in perturbation theory, if the VEVs in Eq.(4.10) are also given as a perturbative expansion (calculated in CPTh) [Bu 93].

First of all this result is very useful in practice, because, given the original interaction, it allows one immediately to write down an ansatz for the effective LF interaction — even if the VEVs cannot, in general, be calculated from the LF Hamiltonian. Secondly, although derived perturbatively,

Eq.(4.10) formally agrees with Eq.(3.29), which was derived nonperturbatively using ε-coordinates. Of course, while Eq.(4.10) was derived only for cases where the VEVs can be calculated perturbatively, Eq.(3.29) is valid in general. However, the formal agreement between the two results gives us confidence to approach other zero-mode problems using perturbation theory as well.

As it stands, Eq.(4.10) is valid only for superrenormalizable theories because we have only addressed longitudinal divergences in the above discussion. For renormalizable theories one must first cut off the transverse divergences, e.g. by using a transverse lattice (see Section 5.3) or dimensional regularization in the transverse direction [MP+ 91, BL 91a]. However, with such a transverse cutoff in place, Eq.(4.10) is valid for field theories in more than one spatial dimension as well.

4.2 Fermions

For the applications of LF quantization to DIS, we are of course not interested in self-interacting scalar fields but rather in theories with fermions and gauge fields. As a first step towards this direction, let us consider fermions interacting with pseudoscalar mesons via a Yukawa coupling (see also Section 2.3.2)

$$\mathcal{L}_{int} = g_p \bar{\psi} i\gamma_5 \psi \chi \tag{4.11}$$

within the framework of LFPTh. First one may be tempted to expect that abovementioned perturbative zero-mode problem does not occur here, because *a priori* there are no tadpoles in Yukawa theory with γ_5-coupling. However, after eliminating the non-dynamical component of the fermion field ($\psi_{(-)}$) from the theory, the canonical LF-Hamiltonian (2.28) does contain terms which are fourth order in the fields — giving rise so-called "seagull"-diagrams (Fig.4.4). It is thus not very surprising that the perturbative zero-mode problem arises in diagrams which have the topology of a seagull with one vacuum contraction. (Figure 4.5), because this is the topology one obtains if one replaces either $\psi^\dagger_{(+)}\partial_-^{-1}\psi_{(+)}$ or ϕ^2 by their VEVs. In practice, this works out as follows [BL 91a, BL 91b, LB 93]: consider, for example, the dressed one loop self-energy diagram for a fermion [4]

$$\Sigma(p) = g_P^2 \int \frac{d^D k}{(2\pi)^D} \int_0^\infty d\mu^2 \frac{\rho(\mu^2)}{(p-k)^2 - \mu^2 + i\varepsilon} \int_0^\infty dm^2 i\gamma_5 \frac{\not{k}\rho_1(m^2) + \rho_2(m^2)}{k^2 - m^2 + i\varepsilon} i\gamma_5, \tag{4.12}$$

[4]Here we assume self-consistently that all sub-loop counterterms have been added to the LF result, such that the full fermion propagator is covariant.



Figure 4.4: Seagull diagram in x^+ ordered perturbation theory, representing four-point interactions induced by eliminating $\psi_{(-)}$. The dashed lines are bosons and the full lines represent fermions. The "slashed" fermion line corresponds to instantaneous (with respect to LF-time) fermion exchange.

Figure 4.5: Typical self-energy diagrams in Yukawa theory, which have the same topology as a seagull with one contraction. The blob stands for arbitrary self energy insertions.

where the spectral functions (ρ, ρ_1, ρ_2) parameterize the (unspecified) self-energy insertions. They satisfy (follows from the canonical commutation relations) $\int_0^\infty d\mu^2 \rho(\mu^2) = \int_0^\infty dm^2 \rho_1(m^2) = 1$. As far as the k_+ integral is concerned, the most singular term in Σ is the one proportional to $\gamma^+ k_+$. We thus consider [5]

$$\begin{aligned} \Sigma^+ &= \frac{tr(\Sigma\gamma^-)}{4} \\ &= g_P^2 \int \frac{d^D k}{(2\pi)^D} \int_0^\infty d\mu^2 \frac{\rho(\mu^2)}{(p-k)^2 - \mu^2 + i\varepsilon} \int_0^\infty dm^2 \frac{k_+ \rho_1(m^2)}{k^2 - m^2 + i\varepsilon}. \end{aligned} \tag{4.13}$$

To identify the troublemaker we eliminate k_+ in the numerator using the algebraic identity

$$\begin{aligned} \frac{k_+}{(k^2 - m^2 + i\varepsilon)\left((p-k)^2 - \mu^2 + i\varepsilon\right)} = \\ \frac{1}{2p_-} \frac{\left[2(p_- - k_-)p_+ - \vec{p}_\perp^{\,2} + 2\vec{p}_\perp \cdot \vec{k}_\perp + m^2 - \mu^2\right]}{(k^2 - m^2 + i\varepsilon)\left((p-k)^2 - \mu^2 + i\varepsilon\right)} \\ - \frac{1}{2p_-} \left[\frac{1}{((p-k)^2 - \mu^2 + i\varepsilon)} - \frac{1}{(k^2 - m^2 + i\varepsilon)} \right]. \end{aligned} \tag{4.14}$$

The important point here is that the last two terms in Eq. (4.14) give δ-functions in $p_- - k_-$ and k_- respectively after the k_+ integration. These δ-functions are missed in the naive LF Hamiltonian without zero-modes (this is very similar to the tadpoles in self-interacting scalar fields). One finds (we subtract here the one-loop result because this allows to drop the surface term in the complex k_- plane; $\rho^{free}(\mu^2) = \delta(\mu^2 - \mu_0^2)$, $\rho_1^{free}(m^2) = \delta(m^2 - m_0^2)$)

$$\begin{aligned} \left(\Sigma^+ - \Sigma^+_{1-loop}\right)_{covariant} &= \left(\Sigma^+ - \Sigma^+_{1-loop}\right)_{canonical\ LF} \\ &+ \frac{g_P^2}{2p_-} \int \frac{d^D k}{(2\pi)^D} \int_0^\infty d\mu^2 \frac{\rho(\mu^2) - \rho^{free}(\mu^2)}{k^2 - \mu^2 + i\varepsilon} \\ &- \frac{g_P^2}{2p_-} \int \frac{d^D k}{(2\pi)^D} \int_0^\infty dm^2 \frac{\rho_1(m^2) - \rho^{free}(m^2)}{k^2 - m^2 + i\varepsilon}. \end{aligned} \tag{4.15}$$

Since the other component of Σ have no problems from zero-modes this immediately implies

$$(\Sigma - \Sigma_{1-loop})_{covariant} = (\Sigma - \Sigma_{1-loop})_{canonical\ LF}$$

[5] A more detailed study shows that the other components are free of trouble [BL 91a, BL 91b].

$$+\frac{g_P^2\gamma^+}{2p_-}\int\frac{d^Dk}{(2\pi)^4}\int_0^\infty d\mu^2\frac{\rho(\mu^2)-\rho^{free}(\mu^2)}{k^2-\mu^2+i\varepsilon}$$
$$-\frac{g_P^2\gamma^+}{2p_-}\int\frac{d^Dk}{(2\pi)^4}\int_0^\infty dm^2\frac{\rho_1(m^2)-\rho_1^{free}(m^2)}{k^2-m^2+i\varepsilon}. \quad (4.16)$$

This result is very interesting for the following reasons:

- canonical LF quantization disagrees with covariant perturbation theory
- the mistake of canonical LF quantization can be compensated by a counterterm to the mass term in the kinetic energy (but not the mass term appearing in the vertex)
- if one adds the wrong counterterm, rotational invariance and parity invariance for physical observables are broken. This can be used as a renormalization condition to "fine-tune" the coefficient of the counterterm.
- the counterterm is related in a simple way to the spectral function of fermions and bosons which are numerically calculable in a canonical LF-calculation!
- The boson contribution in Eq.(4.16) can even be expressed in terms of a local VEV: $\delta\Sigma^{boson} = \langle 0|:\phi^2:|0\rangle g_P^2\gamma^+/2p_-$. Unfortunately this is not possible for the term containing the fermionic spectral density, which would read $\delta\Sigma^{fermion} = \langle 0|:\bar{\psi}\frac{\gamma^+}{i\partial_-}\psi:|0\rangle g_P^2\gamma^+/2p_-$.

Note that, in order to obtain the full counterterm necessary to establish agreement between a covariant calculation and a canonical LF calculation, one still has to add the one loop counterterm — but this should be obvious and can be easily done. Similar statements hold for fermion loops in the boson self energy. The only difference to the above example is that the difference between a covariant calculation and a canonical LF calculation results in a difference in the bare boson mass; i.e. no space time symmetries can be used to fine-tune the counterterm. However, the difference can still be related to the spectral density of the fermions. Besides the "contracted seagulls", only disconnected vacuum diagrams — which are irrelevant for the dynamics of physical states — suffer from the zero-mode problem. It is thus also sufficient to tune the vertex mass and the kinetic mass independently and those masses and the boson mass independently from the corresponding coefficients in the covariant Lagrangian in order to recover equivalence between covariant calculations and canonical LF calculations. Note however,

that (like in the self-interacting scalar theory) all this holds only *after* rendering the transverse momentum integrals finite (e.g. by means of dimensional regularization [MP+ 91, BL 91a] or a transverse lattice (Section 5.3).

It should also be noted that perturbative zero-modes also play a role in higher-twist parton distributions. There they can lead to violations of naive sum rules as discussed in Refs. [Bu 92b, Bu 92c, Bu 95].

4.3 Gauge Theories

In gauge theories the situation is much less clear than in scalar field theories or Yukawa theories, because of notorious infrared singularities in the LF gauge $A^+ = 0$. Certain attempts have been made to perturbatively renormalize LF-QED [MP+ 91, BL 91a, LB 93] and QCD [HZ 93].

In the context of calculations of the electron's anomalous magnetic moment in QED it has been shown (up to three loops in Feynman gauge and up to two loops in LF gauge) that all $k_- \to 0$ singularities in LFPTh cancel — provided one adds up all diagrams that contribute to a given order in the coupling constant [LB 93]. The regulators used were Pauli-Villars regulators or dimensional regularization in the transverse direction. Furthermore only two extra [6] counterterms are necessary to render the theory UV-finite: a kinetic mass counterterm for the electron (similar to the one discussed in Section 4.2) and a mass term for the transverse photon field. The numerical result for $(g-2)$ thus obtained agrees with the known result from covariant calculations.

Perturbative LF calculations of vertex functions, which employ a Tamm-Dancoff truncation were done in Ref. [MP+ 91] for QED and in Ref. [HZ 93] for QCD. Due to incomplete cancelations of $k_- \to 0$ singularities in the Tamm-Dancoff approximation, infrared singular counterterm functions were already in lowest nontrivial order necessary to render the results finite.

The first calculation, relevant for asymptotic freedom, was performed in Ref. [Th 79] (four gluon vertex) and Ref. [CF+ 80] (quark-gluon vertex). Further discussions on renormalization on QCD in LF gauge (but not LF quantization) can be found in Ref. [BD+ 85]. For demonstrations of asymtotic freedom, employing both LF gauge and LF quantization, see Ref. [Pe 94b] (and references therein).

In both types of calculation (LFPTh and LFTD) even perturbatively the structure of the renormalized LF-Hamiltonian is not known to higher orders. Perhaps the cleanest way to address the problem would be to start from the axial gauge in ε-coordinates in a finite box [LN+ 94a, LN+ 94b, LN+ 94c]

[6] That is beyond those counterterms which are required in a covariant calculation.

and to approach the LF by carefully taking the limit $\varepsilon \to 0$ (as in Section 3.3). While it is conceivable that this is feasible in QED, the axial gauge Hamiltonian for QCD_{3+1} in a finite box [LN+ 94c] is perhaps too complicated to allow one to study this limit with appropriate care.

4.4 Summary

In the renormalization of LF field theory, one can distinguish three kinds of counterterms. First the usual renormalizations, which can be handled by making the bare coupling constants in the Lagrangian cutoff dependent. In the following these will be referred to as canonical counterterms. Secondly, counterterms that have to be added when one is employing a Tamm-Dancoff cutoff. These will be discussed in Section 5.4. Typically, one needs an *infinite* number of counterterms! Third, effects caused by an improper treatment of zero-modes in the canonical approach. In those cases were these effects are now understood the renormalization of zero-mode effects can be accomplished by adding a *finite* number of counterterms that have the structure of tadpole and seagull diagrams with some lines "ending in the vacuum". In general, the zero-mode counterterms are already included in the list of "Tamm-Dancoff approximation counterterms". This means zero-mode effects become irrelevant when one uses a Tamm-Dancoff approximation. However, in the absence of a Tamm-Dancoff approximation, i.e. in calculations without or with negligible restrictions on the Fock space (see Section 5.3) or in perturbation theory if one adds all diagrams to a given order in the coupling, abovementioned tadpole or seagull counterterms are quite relevant because they are the only counterterms needed besides the canonical counterterms.

Chapter 5

Nonperturbative Calculations

5.1 Discrete Light-Cone Quantization

The most straightforward method for solving bound state problems in the context of LF quantization is discrete light-cone quantization [1] (DLCQ) [PB 85]. For extensive reviews and more references see Refs. [BP 91, Pa 93, BM+ 93].

The basic idea in DLCQ is as follows (for simplicity we illustrate the method using the example of ϕ^4_{1+1}). One puts the system into an x^--box of length L with periodic or antiperiodic boundary conditions [2]

$$\phi(x^- + L, x^+) = \pm\phi(x^-, x^+). \tag{5.1}$$

In the following, antiperiodic boundary conditions will be used, which implies for the mode expansion

$$\phi(x^-) = \frac{1}{\sqrt{4\pi}} \sum_{k=1}^{\infty} \frac{\left[a_k e^{-ip^k_- x^-} + a_k^\dagger e^{ip^k_- x^-}\right]}{\sqrt{k - \frac{1}{2}}}, \tag{5.2}$$

where

$$p^k_- = \frac{2\pi}{L}\left(k - \frac{1}{2}\right). \tag{5.3}$$

The main reason for choosing antiperiodic boundary conditions is that one does not have to worry about the mode with $p_- = 0$. Another reason

[1] Like the canonical quantization discussed in Chapter 2, the quantization surface in DLCQ is the plane $x^+ = 0$, i.e. a front or plane — and not a cone. Thus discrete light *front* quantization (DLFQ) would be a more appropriate terminology. However, because of historical reasons, the method has been named DLCQ in the literature.

[2] In the presence of interactions which contain odd powers of ϕ one has no choice and one must use periodic boundary conditions — otherwise momentum conservation is violated at the boundary!

is that many numerical problems converge faster when antiperiodic boundary conditions are used (compared to periodic boundary conditions with the $p_- = 0$ mode left out). This can be understood in perturbation theory because there are often non-negligible contributions to Feynman integrals from the region near $p_- = 0$. Let f be some typical function that appears as the argument of some Feynman integral. Then $\varepsilon \sum_{n=-\infty}^{\infty} f\left((n-\frac{1}{2})\varepsilon\right)$ is usually a better approximation to $\int_{-\infty}^{\infty} f(x)$ than $\varepsilon \sum_{n=-\infty}^{-1} f(n\varepsilon) + \varepsilon \sum_{n=1}^{\infty} f(n\varepsilon)$ because in the latter expression the point $n = 0$ is missing compared to the trapezoidal quadrature formula.
In order for $\phi(x)$ to satisfy the canonical commutation relations (see Chapter 2),

$$[\partial_- \phi(x), \phi(y)]_{x^+=y^+} = -\frac{i}{2}\delta(x^- - y^-) \tag{5.4}$$

we impose the usual commutation relations for the coefficients a_k,

$$\left[a_k, a_q^\dagger\right] = \delta_{kq}. \tag{5.5}$$

The above expansion is then inserted into the momentum operator

$$\begin{aligned} P_- &= \int_0^L dx^- : \partial_- \phi \partial_- \phi : \\ &= \frac{2\pi}{L} \sum_{k=1}^{\infty} a_k^\dagger a_k \left(k - \frac{1}{2}\right) \end{aligned} \tag{5.6}$$

and the Hamiltonian

$$\begin{aligned} P_+ &= \int_0^L dx^- \frac{m^2}{2} : \phi^2 : + \frac{\lambda}{4!} : \phi^4 : \\ &= \frac{L}{2\pi}\left(T + V\right), \end{aligned} \tag{5.7}$$

where

$$T = \frac{m^2}{2} \sum_{k=1}^{\infty} \frac{a_k^\dagger a_k}{k - \frac{1}{2}} \tag{5.8}$$

is the kinetic term and

$$V = \frac{\lambda \delta_{P_f P_i}}{8\pi 4!} \sum_{k_1,k_2,k_3,k_4=1}^{\infty} \frac{: \left(a_{k_1}^\dagger + a_{k_1}\right)}{\sqrt{k_1 - \frac{1}{2}}} \frac{\left(a_{k_2}^\dagger + a_{k_2}\right)}{\sqrt{k_2 - \frac{1}{2}}} \frac{\left(a_{k_3}^\dagger + a_{k_3}\right)}{\sqrt{k_3 - \frac{1}{2}}} \frac{\left(a_{k_4}^\dagger + a_{k_4}\right) :}{\sqrt{k_4 - \frac{1}{2}}} \tag{5.9}$$

is the interaction term. $\delta_{P_f P_i}$ is a momentum conserving Kronecker δ. Since the length of the box completely factorizes, it is useful to work with the

rescaled operators

$$K = P_- \frac{L}{2\pi} \tag{5.10}$$

$$H = P_+ \frac{2\pi}{L}. \tag{5.11}$$

Since the momenta of all excitations are discrete and positive, the Fock space is finite dimensional for all K. Thus, at least in principle, one can now proceed as follows: for fixed K (K and H commute) one diagonalizes H (which is a finite matrix for finite K). From the eigenvalues E_i one computes the invariant masses $M_i^2 = 2KE_i$ and from the eigenstates one can compute other physical observables (like parton distributions). In general, physical observables thus computed will of course depend on the "resolution" K. The continuum limit is obtained by extrapolating to $K \to \infty$. The diagonalization is generally done using brute force matrix diagonalization or, if one is only interested in the lowest states, using the Lanczos algorithm [Hi 91].

At this point one encounters a problem that is inherent to Hamiltonian systems: *the dimension of multi-particle states in the Fock space expansion grows exponentially with the number of particles.* The number of particles, as well as the number of states for a single particle are both limited by the longitudinal momentum K, i.e. the dimension of the Fock space basis shows factorial growth with K. Fortunately, in $1+1$ dimensional examples, the factorial growth sets in only rather slowly and numerical convergence for typical observables can be obtained before the size of the matrices becomes a problem. DLCQ was enormously successful in many $1+1$ dimensional field theories [EP+ 87, HV 87, EP 89, Mc 88, Bu 89b, HP+ 90, Bu 89a, Bu 93, Bu 89c, BB 91]. In all cases, where results from other approaches to field theories were available agreement could be shown within numerical uncertainties (QED_{1+1}: [Co 76] vs. [EP+ 87, EP 89] [3] ,QCD_{1+1} [Ha 82] vs. [HP+ 90, Bu 89a], sine-Gordon model: [DH+ 75] vs: [Bu 93]). Beyond reproducing known results, DLCQ has been used to calculate new and interesting results in QCD_{1+1}: the most notable results are the existence of a nucleon-nucleon bound state and the analysis of the nuclear quark distribution in comparison with the nucleon quark distribution. Not only exhibits the $1+1$ dimensional "deuteron" an EMC-effect, but it can also be understood analytically due to the simplified dynamics in $1+1$ dimensions [Bu 89b]. Typical Euclidean lattice calculations are too "noisy" to even

[3] However, there is still a 1% difference in the fundamental meson mass for the term linear in m_e in QED_{1+1} as calculated from bosonization [Co 76] and in LF-quantization [Be 77]. It is not clear whether this deviation is due to the finite Fock space truncation or whether this is a real problem [Ma 92, St 92].

demonstrate binding of hadrons. Another remarkable result from DLCQ calculations in QCD_{1+1} dimensions is "Anti-Pauli-Blocking" [Bu 89c, BB 91]: contrary to the naive expectation, sea quarks in nucleons in QCD_{1+1} tend to have the same flavor as the majority flavor among the valence quarks (i.e. more $\bar{u}$ than $\bar{d}$ in a nucleon $\psi_{valence} = uud$).

In $2+1$ or $3+1$ dimensions the situation changes drastically, because there the exponential growth is much more rapid. The basic reason is that there are now transverse degrees of freedom besides the longitudinal degrees of freedom. Suppose that each particle can occupy N states for each spatial dimension. Then the Fock space basis size grows like $N^{3N_{part}}$ with the number of particles in $3+1$ dimensions, while the corresponding growth would be only $N^{N_{part}}$ in $1+1$ dimensions. For a concrete example (ϕ^4 with antiperiodic boundary conditions in the longitudinal direction) this works out as follows. For a longitudinal momentum $K = \frac{15}{2}$ (8 longitudinal momentum states accessible) the Fock space basis size is 27. If one has just two transverse degrees of freedom (e.g. two points in the transverse direction) the basis size grows to 426. For $8 \times 8 = 64$ degrees of freedom in the transverse direction, that number grows to $6 \cdot 10^{15}$. These astronomical numbers clearly demonstrate that any direct matrix diagonalization approach or even a Lanczos type algorithm is doomed to fail because one is not even able to store the wavefunction in a computer [Bu 94a].

The most simple (and perhaps most drastic) way out of this dilemma is to impose additional cutoffs, like restricting the number of particles. Typically, this means restricting the Fock space to 3 (perhaps 4) particles or less [KP 92, Wo 93]. In QED, since the coupling is small, this is a good approximation. However, to the same order in α within which the 3 particle truncation is a good approximation one can calculate the parton distributions analytically [Bu 92a]. That is, even in QED there is not much point in doing numerical DLCQ calculations with Fock space truncations to the lowest nontrivial order! In QCD, where one faces an intrinsically strong coupling problem, restricting the Fock space to the lowest nontrivial component seems entirely useless. For example, even if one allows up to 4 particles (which is about the maximum that can be handled numerically using the Lanczos algorithm), this means one allows at most one gluon in addition to the three valence quarks in a proton. That is there is no chance one can "see" any effects from nonlinear gluon-gluon couplings. [4]

It should be emphasized, that this problem is not specific for LF field theories, but occurs in many Hamiltonian approaches to field theory — and in many cases could be solved. Thus there are many numerical methods avail-

[4] A caveat to this pessimistic point of view will be discussed in Section 5.4.

able which can potentially be useful in overcoming the difficulties associated with exponential basis size growth.

5.2 Functional Integration on a Longitudinal Lattice

Functional integrals on Euclidean lattices have been very successful in solving ground state properties of QCD (e.g. vacuum properties, hadron masses and ground state matrix elements). However, since two points on a Euclidean lattice are always separated by a space-like distance, it is only very indirectly possible to extract information about light-cone correlation functions from these calculations. Of course, this is because in conventional Euclidean field theory $\exp(-\beta P^0_E)$ is used to project on the ground state wave function of P^0 at equal time. Thus as a caveat one might be tempted to consider a similar formalism for LF Hamiltonians. Suppose one discretizes the x^- direction, [5] and uses a functional integral to project on the ground state of P_+ [Mu 88, HL 93]. This results in an immediate problem because the LF-energy *decreases* with increasing momentum ($P_+ = M^2/2P_-$ in the continuum, on a lattice there is a minimum for $P_- = \mathcal{O}(1/a)$). Due to Bragg reflections, momentum is not conserved and the particles tend to accumulate near the minimum. However, since the momentum near that minimum is of the order of the inverse lattice spacing, the particles always "see" the lattice and no meaningful continuum limit is obtained. It is conceivable that this problem can be cured by adding a Lagrange multiplier proportional to the total LF momentum to the lattice action (in the continuum limit this amounts to minimizing $\tilde{P}_+ = P_+ + \lambda P_-$ instead of P_+). However, this idea will not be investigated here any further. Another difficulty of the longitudinal LF-lattice is species doubling for bosons [Mu 88]!

5.3 Hamiltonian Monte Carlo on a Transverse Lattice

While Mont Carlo calculations for longitudinal LF-lattices seem to be plagued with difficulties, this is not the case for the transverse lattice [Bu 94a]. On a transverse lattice one keeps the longitudinal directions (x^+ and x^-) continuous, while discretizing the transverse coordinate (Fig. 5.1) [BP 76] For simplicity, let us consider self-interacting scalar fields in 2+1

[5] The transverse coordinates are irrelevant in this argument.



Figure 5.1: Space time view of a transverse lattice

dimensions on such a transverse lattice characterized by the action

$$A^{cont.} = \int d^3x \left[\sum_{\mu=0}^{2} \frac{\partial_\mu \phi \partial^\mu \phi}{2} - \frac{m^2}{2}\phi^2 - \mathcal{L}_{int}(\phi) \right]. \tag{5.12}$$

Upon discretizing the transverse direction (spacing a) one thus obtains

$$A^{\perp\ latt.} = a \sum_n \int d^2x \left[\sum_{\mu=0}^{1} \frac{\partial_\mu \phi_n \partial^\mu \phi_n}{2} - \frac{(\phi_{n+1} - \phi_n)^2}{2a^2} - \frac{m^2}{2}\phi_n^2 - \mathcal{L}_{int}(\phi_n) \right]. \tag{5.13}$$

Up to a factor of a (which can be absorbed into a redefinition of the field ϕ_n), Eq.(5.13) looks like the action for a multi-flavor theory in $1+1$ dimensions (where the site index n corresponds to the "flavor" index). In the next step one constructs the DLCQ Hamiltonian for this "multi-flavor" $1+1$ dimensional theory. The important point here is that the action is local in the transverse direction, i.e., there are only nearest neighbor interactions. Since the DLCQ-Hamiltonian is thus also local,

$$H_{DLCQ} = \sum_n \left[H_n + V_{n,n+1} \right], \tag{5.14}$$

one can apply many Monte Carlo techniques which have been developed for other Hamiltonian systems (see e.g. Ref. [NO 87]). One technique which turns out to be particularly useful for LF-Hamiltonians on a transverse lattice is the ensemble projector Monte Carlo technique [DP 85] based on the so called checkerboard decomposition of the Hamiltonian [HS+ 82]. Using

locality of the Hamiltonian one can write

$$H_{DLCQ} = H_a + H_b \tag{5.15}$$

where

$$\begin{aligned} H_a &= \sum_{n=1,2,3,...} \frac{H_n}{2} + \sum_{n=1,3,5,...} V_{n,n+1} = H_{1,2} + H_{3,4} + ... \\ H_b &= \sum_{n=1,2,3,...} \frac{H_n}{2} + \sum_{n=2,4,6,...} V_{n,n+1} = H_{2,3} + H_{4,5} + ... \end{aligned} \tag{5.16}$$

In other words, the DLCQ Hamiltonian can be written as a sum of two terms, each of which can be written as a direct sum of two-site-Hamiltonians. The point to all this is that while the dimension of the space on which H_{DLCQ} acts is astronomical, the two-site-Hamiltonians act only on a very small Hilbert space (for our above example with $K = \frac{15}{2}$ and say 16 transverse sites: $dim(H_{DLCQ}) = 7.8 \cdot 10^8$ but $dim(H_{n,n+1}) = 426$). The method is called checkerboard algorithm because one approximates the time evolution operator of the system by alternating infinitesimal time evolution operators generated by H_a and H_b respectively

$$e^{-\varepsilon H_{DLCQ}} = e^{-\frac{\varepsilon}{2}H_a} e^{-\varepsilon H_b} e^{-\frac{\varepsilon}{2}H_a} + \mathcal{O}(\varepsilon^3). \tag{5.17}$$

If one axis of the checkerboard is the discretized space direction and the other the time, Eq. (5.16) can be interpreted as if interactions between sites occur only across the black squares [HS+ 82].

Before we explain how the infinitesimal time evolution operators are multiplied together, let us pause here for a moment and understand the advantage of the transverse lattice with DLCQ over the longitudinal lattice discussed in the previous Section. The main cause of the problem in the previous Section was lack of longitudinal momentum conservation. In DLCQ P_-, the longitudinal momentum, is manifestly conserved. Furthermore, P_- is just the sum of momenta at each site

$$P_- = \sum_n P_-^n, \tag{5.18}$$

and the checkerboard algorithm is compatible with with longitudinal momentum conservation ($[H_a, P_-] = [H_b, P_-] = 0$). I.e. with DLCQ on a transverse lattice, longitudinal momentum is conserved at each step of the calculation. Hence, one can minimize P_+ while keeping P_- manifestly fixed and there are no "runaway solutions".

In the actual calculations one uses Monte Carlo techniques to calculate $\left(e^{-\varepsilon H}\right)^N |\psi_i(K)\rangle$ by alternate application of $e^{-\frac{\varepsilon}{2}H_a}$, $e^{-\varepsilon H_b}$, $e^{-\varepsilon H_a}$,...,$e^{-\frac{\varepsilon}{2}H_a}$ to

$|\psi_i(K)\rangle$. Here $|\psi_i(K)\rangle$ is an initial guess for the ground state wave function with longitudinal momentum K. For $N \to \infty$ one thus obtains an approximation (because ε is finite, the result is not exact) to the ground state hadron with the same good [6] quantum numbers as $|\psi_i(K)\rangle$. One very useful technique is the ensemble projector Monte Carlo method [NO 87, DP 85], which works as follows for these systems:

1) let $|n\rangle$ be a complete set of states
(here product basis of Fock state bases at each site)

2) make a good guess for $|\psi_i(K)\rangle$
(here a valence state with $P_\perp = 0$: $|\psi_i(K)\rangle = \sum_{n_\perp} a^\dagger_{K,n_\perp}|0\rangle$)

3) start from ensemble $|i^{(0)}_\nu\rangle$ of states (from set $|n\rangle$)

4) for each $|i^{(0)}_\nu\rangle$ select a new state $|i^{(1)}_\nu\rangle$ with probability

$$W(i^{(1)}_\nu, i^{(0)}_\nu) = \frac{|\langle i^{(1)}_\nu|e^{-\frac{\varepsilon}{2}H_a}|i^{(0)}_\nu\rangle|}{\sum_n |\langle n|e^{-\frac{\varepsilon}{2}H_a}|i^{(0)}_\nu\rangle|}$$

and calculate the score

$$S^{(1,0)}_\nu = \frac{\langle i^{(1)}_\nu|e^{-\frac{\varepsilon}{2}H_a}|i^{(0)}_\nu\rangle}{W(i^{(1)}_\nu, i^{(0)}_\nu)}$$

note: $W(i^{(1)}_\nu, i^{(0)}_\nu)$ factorizes into two-site probabilities
$\hookrightarrow$ local (one pair of sites at a time) "updating" possible

5) replicate states with multiplicity:
$int\left[\frac{|S^{(k,k-1)}_\nu|}{S} + \text{random number} \in (0;1)\right]$

($\bar{S}$: av. score)

Thus paths with large scores contribute with multiple weight, while paths with small scores get eliminated.

6) repeat while alternating H_a and H_b
($\frac{1}{2}$ in exponent only in 1^{st} and last step!)

[6] Of course, only those quantum numbers which are associated with exact symmetries of DLCQ on a transverse lattice (like C-parity or baryon number in QCD) are relevant here.

7) observables from ensemble average, e.g., energy of ground state hadron:

$$E_0 = \lim_{N\to\infty} \frac{\sum_\nu \langle\psi_f(K)|H|i_\nu^{(N)}\rangle sign\left[S_\nu^{(N,N-1)}\cdot ...\cdot S_\nu^{(1,0)}\right]}{\sum_\nu \langle\psi_f(K)|i_\nu^{(N)}\rangle sign\left[S_\nu^{(N,N-1)}\cdot ...\cdot S_\nu^{(1,0)}\right]}$$

other observables (e.g. an observable diagonal in the basis)

$\langle\psi_0(K)|\hat{O}|\psi_0(K)\rangle =$

$$\lim_{M,N\to\infty} \frac{\sum_\nu \langle\psi_f(K)|i_\nu^{(N+M)}\rangle sign\left[S_\nu^{(N+M,N+M-1)}\cdot ...\cdot S_\nu^{(1,0)}\right]\langle i_\nu^{(N)}|\hat{O}|i_\nu^{(N)}\rangle}{\sum_\nu \langle\psi_f(K)|i_\nu^{(N+M)}\rangle sign\left[S_\nu^{(N+M,N+M-1)}\cdot ...\cdot S_\nu^{(1,0)}\right]}$$

In this Monte Carlo procedure, one only has to store the ensemble of states at one "timeslice" plus the result of the measurement of the observable after N slices. Thus, at least in principle, one can handle very large lattices. The main advantages of the transverse lattice are as follows [Bu 94a]

- longitudinal momentum is manifestly conserved → no runaway solutions
- parton distributions are diagonal in the DLCQ-basis
- LF-vacuum is trivial → no statistical fluctuations from updating the vacuum far away from physical states on huge lattices.
- species doubling for fermions occurs only for the latticized transverse dimensions → can be easily compensated by staggering [Gr 93].
- excited states are suppressed by the square of their masses:
  $\exp(-N\varepsilon P_+) = \sum_n |n\rangle \exp(-N\varepsilon M_n^2/2P_-)\langle n|$ instead of
  $\exp(-N\varepsilon P^0) = \sum_n |n\rangle \exp(-N\varepsilon M_n)\langle n|$ which one encounters in a conventional Hamiltonian formulation.

It is interesting to see how confinement emerges on the transverse lattice in the limit of large lattice spacing: In this limit, the coupling between the sheets is weak and the energy scale associated with link field excitations is high. Therefore, when one separates two test charges in the longitudinal direction, the transverse lattice behaves similar to QCD_{1+1} and linear confinement results trivially. For transverse separations between the charges, a different mechanism is at work. Gauge invariance demands that the two charges are connected by a string of link fields. In the limit of large spacing the link fields fluctuate only little and the energy of such a configuration can

be estimated by counting the number of link fields needed to connect the charges, which again yields linear confinement.

Some of the disadvantages of the transverse lattice are: Since $x^+ \to ix^+$ is *not* a Wick rotation (it is just a mathematical trick to project on the ground state of P_+), the metric is not Euclidean and thus propagators oscillate. Hence, negative scores occur already for bosons which leads to an increase in the statistical fluctuations. However, these negative scores turn out to have only a small statistical weight and the resulting "sign-problem" is not serious. Very often in LF calculations, large cancellations occur between different terms in the Hamiltonian. For example, the instantaneous photon exchange has a $1/q_-^2$ singularity which is canceled by vertex factors in photon exchange. In general, it is difficult to obtain such cancellations from a Monte Carlo calculation. Another difficulty is that gauge invariance on a lattice can only be maintained if one introduces link fields. On a transverse lattice this amounts to introducing $1+1$-dimensional gauged nonlinear sigma model fields on each link [BP 76]. Constructing a Fock space basis out of these nonlinear degrees of freedom and calculating appropriate matrix elements is a nontrivial task [Gr 92b, Gr 94b].

The sign problem associated with fermions is a notorious difficulty for Monte Carlo algorithms: due to the minus sign in exchange terms, the infinitesimal time evolution operator tends to contain many negative matrix elements. This very general problem is also expected to afflict Mont Carlo calculations on transverse lattices. However, since the LF vacuum is trivial, there are no sign fluctuations from Z-graphs and vacuum diagrams. Thus one expects that the sign problem on the LF is less severe than usual. Whether this improvement is sufficient to render fermions tractable on transverse lattices has not yet been investigated.

Obviously, the transverse lattice lacks manifest rotational invariance which must be restored in the process of renormalization. Recently, a technique has been described that allows easy computation of the potential between infinitely heavy quarks in a LF framework [Bu 94c]. Demanding rotational invariance for this observable may prove to be a powerful tool in such a procedure.

5.4 Light-Front Tamm-Dancoff

As we have discussed already in Section 5.1, the dimensionality of the Fock space grows dramatically as one includes higher Fock components. Clearly, since α_S is fairly large at a low momentum scale, a numerical solution of bound state problems in QCD (which includes all scales) necessarily involves

many Fock components. In this chapter we will discuss the light-front Tamm-Dancoff (LFTD) approach to LF problems (for a comprehensive review see Ref.[WW+ 94]). The basic idea is very simple [PH+ 90, WR 94]: Hadrons are complicated objects only if one tries to build them in terms of bare quarks and gluons whose masses and couplings are renormalized at a scale of 1 GeV or higher. In terms of collective excitations (constituent quarks) ground state hadrons are rather simple. One of the problems with the constituent quark model is that *a priori* the interactions among the quarks are *ad hoc*.

The goal of LFTD field theory is to systematically eliminate higher Fock components and high energy degrees of freedom[7]. As one goes to lower and lower scales the interaction between the (dressed) constituents thus becomes more and more complicated. If the whole program is successful, constituent quarks will emerge as the quasiparticles of QCD at intermediate energies. A major virtue of using LF quantization in this approach is that it stays close both to physical intuition (which may prove very helpful when it comes to developing variational methods to analyse the Hamiltonian) as well as to experimental observables at large momentum transfer (useful for phenomenological applications).

A systematic Fock space expansion, based on a Hamiltonian formulation, for field theory was originally developed by Tamm [Ta 45] and independently by Dancoff [Da 50]. It turns out that such an approach is doomed to fail if the perturbative ground state (the starting point of the expansion) is too far from the actual ground state. In such a situation one needs very (or infinitely) complicated Fock states just to build the ground state. LF quantization is advantageous at this point, because the vacuum of a LF Hamiltonian is trivial.

In fact, because of the vacuum, LF quantization is probably the *only* framework, where such a programm can possibly work.

In practice, even within LF quantization, it is of course not possible to integrate out high energy states and higher Fock states exactly. Instead one writes down a catalog of all interaction terms that are allowed by power counting [Wi 91, GP 92]: on the LF there are two length scales x^- and $x_\perp$. The engineering dimensions of the various operators and terms that enter the LF-Hamiltonian in $3+1$ dimensions can be easily derived from free field theory [GP 92] (ϕ stands for a scalar field or for $A_\perp$ — the transverse gauge field components which have the same engineering dimension as scalar fields;

[7]This procedure is explicitely demonstrated for the simple example of ϕ^4_{3+1} in the two particle sector in Ref. [Am 94].

$\psi^{(+)}$ is the dynamical component of a fermion field)

$$\partial_- = \left[\frac{1}{x^-}\right],\ \partial_\perp = \left[\frac{1}{x_\perp}\right],\ m = \left[\frac{1}{x_\perp}\right] \tag{5.19}$$

$$\psi^{(+)} = \left[\frac{1}{x_\perp\sqrt{x^-}}\right],\ \phi = \left[\frac{1}{x_\perp}\right]. \tag{5.20}$$

The Hamiltonian and the Hamiltonian density have dimensions

$$H = \left[\frac{x^-}{x_\perp^2}\right],\ \mathcal{H} = \left[\frac{1}{x_\perp^4}\right]. \tag{5.21}$$

Thus all allowed terms without fermion fields are [GP 92] [8]

$$m^3\phi,\ m^2\phi^2,\ \partial_\perp^2\phi^2,\ m\phi^3,\ \phi^4. \tag{5.22}$$

Including fermion fields one obtains [GP 92]

$$\frac{1}{\partial_-}\psi^{(+)\dagger}\Gamma\left\{m^2, \partial_\perp^2, m\vec{\gamma}_\perp\vec{\partial}_\perp, m\phi, \phi^2\right\}\psi^{(+)}, \tag{5.23}$$

$$\frac{1}{\partial_-^2}\psi^{(+)\dagger}\Gamma_1\psi^{(+)}\psi^{(+)\dagger}\Gamma_2\psi^{(+)} \tag{5.24}$$

(Γ, Γ_1 and Γ_2 are some Dirac matrices). Unfortunately, this is not the whole story. Already free LF-field theory is nonlocal in the longitudinal direction [e.g. for scalar fields because of the fundamental commutator $[\phi(x^-,x^+),\phi(y^-,x^+)] = -\frac{i}{4}\varepsilon(x^- - y^-)$ and for fermions because an inverse derivative of ∂_- appears in the kinetic energy term (2.28)]. Thus longitudinal locality is no longer a restriction on the functional form of the possible terms in the Hamiltonian. As a result, any of the operators in the above catalog may be multiplied by arbitrary functions of ratios of longitudinal momenta! In fact, there are examples known where complicated functions of ratios of incoming and outgoing momenta, multiplying a four fermion counterterm, are necessary to cancel UV divergences [GH+ 93].

As a result of this infinitely complicated counterterm structure, it seems one looses predictive power and one thus might be forced to abandon LFTD as a *fundamental* theory of hadrons. It should be emphasized that, even if one does have to abandon LFTD as a fundamental theory, it might still have many virtues in parton phenomenology. However, there has been recent progress towards understanding how to apply renormalization group techniques to LFTD that may help restore its predictive power [WW+ 94] (for an excellent pedagogical review, see Ref. [Pe 94b]).

Unless one works with the full Hamiltonian, nonperturbative bound state calculations in QCD almost inevitably violate gauge invariance [9]. Therefore,

[8] See the discussion in Ref.[GP 92] why terms with negative powers of m are excluded.

[9] Lattice gauge theory being the only exception.

if one wants to derive a constituent picture from QCD, one is forced to allow that explicit gauge invariance is violated: *gauge invariance becomes a hidden symmetry* [WW+ 94, Pe 94b]. In LFTD one introduces cutoffs that violate symmetries which normally prevent a constituent picture from arising (gauge invariance and full Lorentz invariance). In a sense, the counterterm functions that complicate renormalization offer a possible resolution of apparent contradictions between the constituent picture and QCD [Pe 94b].

The technique to remove cutoff dependence from physical results is renormalization: for example, the functions of momentum fractions that appear in the relevant and marginal operators can be fixed by demanding covariance and gauge invariance in physical observables. An alternative way to fix the marginal and relevant counterterms is *coupling constant coherence* (CCC): one insists that functions appearing in non-canonical relevant and marginal operators are not independent functions of the cutoff, but depend on the cutoff implicitly through their dependence on canonical couplings [Pe 94a, Pe 94b]. This automatically fixed they way in which new variable evolve with the cutoff, and it also fixes their value at all cutoffs if one insists that the new counterterms vanish when the canonical couplings are turned of [Pe 94b]. Remarkably, in the examples studied in Ref. [PW 93, Pe 94a], this procedure provided the precise values for the non-canonical terms that were required to restore Lorentz covariance for physical observables. For an explicit example for CCC, the reader is referred to Ref. [Pe 94b].

Even without assuming CCC, one can employ renormalization group techniques [Wi 75] to help determine the counterterm functions: using the powerful tool of try and error, one makes an ansatz for these functions, which one can improve by repeatedly applying renormalization group transformations to the effective LF Hamiltonians with these functions included. The fact that the QCD Hamiltonian should be an ultraviolet stable fixed point under these transformations can be exploited to improve the original ansatz for the counterterm functions [WW+ 94]. Probably, such transformations alone are not sufficient to completely determine the renormalized LF Hamiltonian for QCD, but one can improve this approach considerably by using perturbation theory, CCC and perhaps phenomenology to guide the ansatz functions used in the renormalization group approach to LF Hamiltonians.

Another promising idea in the context of LFTD is the *similarity transformation* [GW 93]. Whenever one derives an effective Hamiltonian by eliminating states above a certain cutoff perturbatively, one faces small energy denominators, and thus large and uncertain corrections, for states that close to (and below) the cutoff. This feature makes it very difficult to repeatedly apply renormalization group transformations because matrix elements of states near the cutoff are large. To resolve this problem, Glazek and Wil-

son have suggested to apply a cutoff to *energy differences* instead of to *single particle energies*. By construction, this resolves the problem of small energy denominators, but it also provides a band diagonal Hamiltonian. The *similarity transformation* exploits this type of cutoff and thus provides a way to apply renormalization group techniques to LF Hamiltonians (and other many body problems) [GW 93].

Chapter 6

Summary, Conclusions and Outlook

LF field theory is a very promising approach toward calculating correlation functions along a light-like direction. Such correlation functions appear in the theoretical analysis of a variety of hard scattering processes, such as deep inelastic lepton-hadron scattering and asymptotic form factors. Probably the most intriguing and controversial property of LF Hamiltonians is the triviality of the ground state. Recent developments indicate that LF Hamiltonians must be regarded as effective Hamiltonians in the sense that some of the interactions acquire nonperturbative renormalizations with coefficients proportional to vacuum condensates. So far one understands the LF vacuum and is able to construct the effective LF-Hamiltonian only in a few toy models. However, in these examples only a finite number of condensates are necessary to completely specify the Hamiltonian. It would be extremely useful if one could construct and approximately solve such an effective LF Hamiltonian for QCD, not only for the analysis of hard processes, but also for our understanding of low energy QCD: due to the triviality of the LF vacuum, a constituent picture makes sense and an effective LF Hamiltonian for QCD would offer the opportunity for deriving a constituent picture as an approximation to the QCD bound state problem.

Three main stream directions can be distinguished in the endeavor toward constructing a LF Hamiltonian for QCD: First, a *fundamental* approach where all zero-modes are included as dynamical degrees of freedom. Second, an *effective* approach where one attempts to absorb all zero-modes and associated vacuum effects into *effective* interactions and coupling constants. Third, the *LF Tamm Dancoff* approach, where not only vacuum effects but also effects from high energy and high Fock components are "integrated out" and absorbed into effective interaction terms. In the *fundamental* approach

(this includes all formulations of LF field theory where explicit zero-mode degrees of freedom are included) the vacuum as well as the physical particle states are complicated and one partly looses the dynamical advantages of the LF framework. While it seems easier to construct the Hamiltonian than in the other two approaches, the main difficulty of the *fundamental* approach lies in the fact that the equations of motion are extremely complicated. It is not clear whether such an approach provides any computational advantage over a conventional Hamiltonian approach. Nevertheless, it is very useful to pursue this approach further in order to provide a solid theoretical basis for other, more practical, approaches to LF field theory. For example, studies that include zero-modes can be useful for deriving an ansatz for the effective LF Hamiltonian in the large volume limit.

The *LF Tamm Dancoff* approach corresponds to the other extreme. The vacuum is trivial and the physical particle states are very simple — by construction they contain only the low energy effective degrees of freedom. A major virtue of this approach is that it stays close to physical intuition and thus potentially offers a connection between the constituent picture and QCD [Pe 94b]. While the *LF Tamm Dancoff* approach is thus very appealing from the intuitive point of view its main disadvantage is the enormous complexity of the effective Tamm Dancoff Hamiltonian. In principle an infinite number of counterterms are possible. These counterterm functions are heavily constrained by imposing Lorentz covariance on physical observables or by demanding cancelation of unphysical divergences. However, so far it is not clear to what extend one can employ renormalization group techniques to constrain the possible interactions to the point where only a few (instead of infinitely many) free parameters enter the LF Tamm Dancoff Hamiltonian of QCD. The second (*effective*, in the sense of zero-mode free) approach toward constructing the LF Hamiltonian for QCD stands in between the other two in several respects. The vacuum is trivial but physical particles will in general have a complicated wavefunction. Some of the interactions in the effective LF Hamiltonian have coefficients proportional to vacuum condensates. Those can either be regarded as free parameters or (in some cases) they can be determined from self-consistency conditions. Surprisingly, in those cases where the construction of such an effective Hamiltonian has been accomplished, already a finite number of condensates is sufficient to specify the Hamiltonian. [1] This is a very encouraging result. Perturbative calcula-

[1]This approach should not be confused with the standard QCD-sum rules approach to the strong interactions [NS+ 81], where one does *not* solve a Hamiltonian and where practically all the dynamics is buried in the condensates. Hence it is not surprising that less condensates are necessary as an input in the LF effective Hamiltonian approach than in the sum rule approach.

tions up to two loops indicate a similar result for QED, where the two loop calculations do not require any counterterms which are not already present at the one loop level. LF perturbation theory in QCD has so far only been performed up to one loop.

Although there has been considerable progress recently, so far none of these three approaches has been successful to the point where it was possible to construct a useful LF Hamiltonian for QCD. The initial optimism about LF quantization, spurred by the very successful application to $1+1$ dimensional field theories, was premature. Much work remains to be done before LF quantization can be applied to QCD.

For example, it is still not completely understood to what extend LF Hamiltonians, with a trivial vacuum, can account for the phenomenon of spontaneous symmetry breaking. The only examples where this subject seems to be mostly understood are ϕ^4 theory in $1+1$ dimensions and field theories in the mean field approximation. It would be interesting to study cases where the order parameter for the symmetry breaking does not enter the Hamiltonian — which is for example the case in the spontaneous breakdown of chiral symmetry in QCD.

A possibly related issue, which requires further study, concerns the *non-covariant counterterms*. In the context of perturbation theory it has been shown that a finite number of such counterterms are necessary in the bare Hamiltonian to recover full Lorentz covariance for physical observables. However, so far it has not been demonstrated that the proposed counterterms are sufficient to restore Lorentz covariance for physical observables in a nonperturbative calculation.

Within the context of LF Tamm-Dancoff it is still necessary to demonstrate that the renormalization group, combined with constraints from Lorentz invariance, is sufficient to fix the infinite number of counterterms which are possible on general grounds.

For the transverse lattice approach to be useful, it must be shown that the fermion sign problem, which usually limits Hamiltonian Monte Carlo calculations with fermions considerably, is tractable. Since vacuum fluctuations are suppressed in LF quantization, any sign problems arising from vacuum diagrams are trivially absent. While this is a very encouraging observation, it resolves only part of the problem — sign problems arising from exchange diagrams within a hadronic state are of course still there. Another difficulty for transverse lattice calculations occurs because gauge invariance on such a lattice requires the introduction of $1+1$-dimensional link fields. One must learn to work with these "nonlinear sigma model" degrees of freedom in the context of LF quantization before one can apply the transverse lattice to QCD.

Besides QCD oriented applications of the LF formalism, it may turn out to be very useful to consider phenomenological and/or more nuclear physics oriented applications as well. For example, it may be interesting to reconsider the pion contribution to nuclear structure functions [FP 83] from the point of view of LF quantization. On the one hand, this could be helpful in clarifying the role of binding effects in such calculations. On the other hand, such works may help to demonstrate the usefulness of LF quantization to people who are not directly involved in the field.

LF quantization is very closely related to the *infinite momentum frame* formulation of field theory. Intuitively one would thus expect that the LF formulation of QCD offers a new theoretical approach to relativistic heavy ion collisions. So far, this connection has been exploited only very little [MV 94].

Acknowledgments

I would like to thank M. Frank for many suggestions that helped to make this article more "readable". I am also very grateful to many colleagues and collaborators for fruitful and enlightening discussions over the last years, particularly with F. Lenz, S. J. Brodsky, A. Langnau, R. J. Perry, E. Swanson and P. Griffin.

Appendix A

The Dirac Bergmann Formalism

In this appendix, a brief introduction into the Dirac-Bergmann quantization procedure [Di 50, Be 56, HR 76] is given. Quite generally, it replaces the canonical quantization procedure in the presence of constraints. However, it can also be used to derive the correct fundamental commutation relations for theories, where the Lagrangian contains at most linear terms in the time derivative. [1]

This is for example the case for many field theories, when expressed in terms of LF-variables [Su 82]. For example, for a noninteracting massive scalar field in $1+1$ dimensions one obtains

$$\mathcal{L} = \partial_+\phi\partial_-\phi - \frac{m^2}{2}\phi^2, \tag{A.1}$$

which contains no terms quadratic in $\frac{\partial}{\partial x^+}$. Naive canonical quantization, i.e.

$$\pi = \frac{\delta\mathcal{L}}{\delta\partial_+\phi} = \partial_-\phi \tag{A.2}$$

$$[\pi(x), \phi(y)]_{x^+=y^+} = -i\delta(x^- - y^-), \tag{A.3}$$

with

$$P_+ = \int dx^- T_{+-} = \int dx^- \frac{m^2}{2}\phi^2 \tag{A.4}$$

yields

$$-\partial_\mu\partial^\mu\phi = -2\partial_+\partial_-\phi = -i\left[P_+, \partial_-\phi\right] = 2m^2\phi. \tag{A.5}$$

[1] It should be noted that, in the latter case, alternate treatments are possible as well [FJ 88].

Clearly, Eq.(A.5) differs from the (correct) Euler-Lagrange equation $-2\partial_+\partial_-\phi = m^2\phi$ by a factor of two [CR+ 73]. This mistake arises because the kinetic term in Eq.(A.1) is only linear in the time derivative ∂_+. Thus the equation relating the canonical momenta to the fields (A.2) is a constraint equation since it contains no time derivative and therefore the phase space variables $\pi(x)$ and $\phi(x)$ for a given time are not independent.

Quantizing a system with constraints is a nontrivial task. Fortunately, the Dirac-Bergmann algorithm provides a step by step prescription for the proper quantization procedure. The basic steps of this procedure will be illustrated in an example below. To keep the discussion simple, zero-modes will be deliberately left out in the discussion. A complete discussion, which includes zero-modes, can be found in Refs.[HK+ 89, HK+ 91b, HK+ 92b, HK+ 91a] Furthermore, the discussion here will be restricted to a system with a finite number of degrees of freedom (which can, for example, be obtained from Eq.(A.1) by discretizing the x^- direction)

$$L(\phi_i, \dot{\phi}_i) = \sum_{i,j=1}^{N} A_{i,j}\dot{\phi}_i\phi_j - \sum_i V(\phi_i) \tag{A.6}$$

with $A_{ij} = -A_{ji}$ (the symmetric part of A_{ij} corresponds to a total time derivative and can be subtracted). The canonical momenta are given by

$$\pi_i = \frac{\partial \mathcal{L}}{\partial \dot{\phi}_i} = \sum_j A_{ij}\phi_j \tag{A.7}$$

with Poisson brackets

$$\begin{aligned} \{\phi_i, \pi_j\} &= \delta_{ij} \\ \{\phi_i, \phi_j\} &= \{\pi_i, \pi_j\} = 0 \end{aligned} \tag{A.8}$$

Eq.(A.7) does not contain any time derivative, i.e. it should be considered as a constraint

$$\chi_i \equiv \pi_i - \sum_j A_{ij}\phi_j \approx 0 \tag{A.9}$$

$i = 1, ..., N$. Here ≈ 0 means *weakly vanishing*, i.e. a constraint on the physical phase space. The *canonical Hamiltonian* is constructed as usual

$$H_c = \sum_i \dot{\phi}_i\pi_i - L = \sum_i V(\phi_i). \tag{A.10}$$

The constraints (A.9) have nonvanishing Poisson brackets with H_c

$$\{\chi_i, H_c\} = -V'(\phi_i) \tag{A.11}$$

as well as among themselves

$$\{\chi_i, \chi_j\} = A_{ij} - A_{ji} = 2A_{ij}. \tag{A.12}$$

Thus if the time evolution would be generated by Poisson brackets with H_c the the theory would be inconsistent because the constraints would not be satisfied at all times. To remedy the situation one adds Lagrangian multipliers to H_c, yielding the *primary Hamiltonian*

$$H_p = H_c + \sum_i \lambda_i \chi_i \tag{A.13}$$

and demands strong vanishing of the Poisson bracket of H_p with the constraints

$$0 = \{\chi_i, H_p\} = \{\chi_i, H_c\} + \sum_k \lambda_k B_{ki} \tag{A.14}$$

where $B_{ki} = 2A_{ki}$. To simplify the discussion, let us suppose that B^{-1} exists. [2] Then one can satisfy Eq.(A.14) by choosing

$$\lambda_k = \sum_i (B^{-1})_{ki} \{\chi_i, H_c\}. \tag{A.15}$$

The primary Hamiltonian thus reads

$$H_p = H_c - \sum_{i,j} \chi_i (B^{-1})_{ij} \{\chi_j, H_c\}. \tag{A.16}$$

Let us now introduce the Dirac brackets between X and Y

$$\{X, Y\}_D = \{X, Y\} - \sum_{i,j} \{X, \chi_i\} (B^{-1})_{ij} \{\chi_j, Y\}. \tag{A.17}$$

By construction one has

$$\dot{X} = \{X, H_c\}_D = \{X, H_p\}. \tag{A.18}$$

Actually, the Dirac bracket of any operator with any of the constraints vanishes identically

$$\begin{aligned} \{X, \chi_k\}_D &= \{X, \chi_k\} - \sum_{i,j} \{X, \chi_i\} (B^{-1})_{ij} \{\chi_j, \chi_k\} \\ &= \{X, \chi_k\} - \sum_{i,j} \{X, \chi_i\} (B^{-1})_{ij} B_{jk} = 0, \end{aligned} \tag{A.19}$$

[2] For the LF-Lagrangian this is actually not the case. There is one zero eigenvalue — the infamous zero-mode — which has to be treated separately. The resulting procedure is known as the modified Dirac-Bergmann algorithm.

i.e. in particular $\{\chi_i, \chi_j\} = 0$. In a sense, the Dirac brackets take all the phase space restrictions from the constraint equations automatically into account. This is in sharp contrast to the Poisson brackets, which are calculated as if all the π_i's and ϕ_i's were independent $\{X, Y\} = \sum_i \frac{\partial X}{\partial \phi_i}\frac{\partial Y}{\partial \pi_i} - \frac{\partial Y}{\partial \phi_i}\frac{\partial X}{\partial \pi_i}$. It thus seems natural to use Dirac brackets, instead of Poisson brackets, when identifying classical brackets with quantum commutators

$$\{X, Y\}_D \to i\,[X, Y] \tag{A.20}$$

in the quantization process. A more thorough discussion on this subject can for example be found in Ref.[Su 82]. Here we are more interested in the consequences of Eq.(A.20). For this purpose, let us evaluate the fundamental Dirac brackets

$$\begin{aligned} \{\phi_i, \pi_j\}_D &= \{\phi_i, \pi_j\} - \sum_{k,l} \{\phi_i, \chi_k\} (B^{-1})_{kl} \{\chi_l, \pi_j\} \\ &= \delta_{i,j} - \sum_{k,l} \delta_{ik} (B^{-1})_{kl} A_{lj} = \frac{1}{2}\delta_{i,j}. \end{aligned} \tag{A.21}$$

Roughly speaking, the reduction of the number of independent degrees of freedom by a factor of two manifests itself in a factor $1/2$ in the Dirac bracket, and after applying Eq.(A.20) the factor $1/2$ also appears in the quantum commutator between π_i and ϕ_j. For the LF-quantization of scalar fields this implies

$$[\pi(x), \phi(y)]_{x^+=y^+} = -\frac{i}{2}\delta(x^- - y^-) \tag{A.22}$$

instead of Eq.(A.3). Clearly, this remedies the abovementioned (A.5) problem with the extra factor of 2 in the LF equation of motion generated by P_+ (A.5).

From the physics point of view [Gr 94a], the whole difficulty in quantization with constraints could be avoided if it were possible to choose degrees of freedom which are compatible with the constraints. For the above example this is actually possible, since the constraint (A.9) is linear in the fields. Let us thus make the ansatz

$$\phi_i^D = \alpha \left[\phi_i + \sum_j \beta_{ij} \pi_j\right], \tag{A.23}$$

where α is a constant and the β_{ij} are determined by requiring a vanishing Poisson bracket between ϕ_i^D and the constraints

$$0 = \left\{\phi_i^D, \chi_j\right\} = \alpha \left[\delta_{ij} + \sum_k \beta_{ik} A_{kj}\right], \tag{A.24}$$

i.e. $\beta_{ij} = (A^{-1})_{ij}$. The normalization α is fixed by demanding that $\phi_i^D \approx \phi_i$, i.e.

$$\phi_i^D = \alpha \left[\phi_i + \sum_j \left(A^{-1} \right)_{ij} \pi_j \right] \approx \alpha \left[\phi_i + \sum_j \left(A^{-1} \right)_{ij} \sum_k A_{jk} \pi_k \right] = 2\alpha \phi_i, \tag{A.25}$$

yielding $\alpha = \frac{1}{2}$. By construction, ϕ_i^D does not "see" the constraint and one can apply canonical quantization rules directly

$$\left[\pi_i, \phi_j^D \right] = -i\delta_{ij} \tag{A.26}$$

and thus

$$[\pi_i, \phi_j] = -i\alpha\delta_{ij} = -\frac{i}{2}\delta_{ij}. \tag{A.27}$$

In the continuum limit this results in Eq.(A.22).

As already indicated above, there are more points that need to be discussed before the Dirac-Bergmann procedure for constructing the LF Hamiltonian is complete. In particular, one has to address the issue of zero-modes. In the LF example, the analog of the matrix A_{ij} is the differential operator ∂_-. Fields which are independent of x^- are annihilated by ∂_- and thus correspond to eigenvectors with eigenvalue zero (on a finite interval, with periodic boundary conditions, these zero-modes have to be considered for a complete formulation of the theory). In such a situation one first has to project on the Hilbert space orthogonal to the zero-modes before the simplified procedure above can be applied. The resulting modified Dirac-Bergmann procedure is quite involved and has been discussed extensively in the literature [HK+ 89, HK+ 91b, HK+ 92b, HK+ 91a]. The basic difficulty arises because the constraint equation for the zero-mode is nonlinear. For example, in ϕ^4 theory in a "box" with periodic boundary conditions in the x^--direction, integrating the Euler Lagrange equation $-2\partial_-\partial_+\phi = m^2\phi + \frac{\lambda}{3!}\phi^3$ over x^- yields [HK+ 92a, PV 93, PV 94]

$$m^2 \int dx^- \phi + \frac{\lambda}{3!} \int dx^- \phi^3 = 0. \tag{A.28}$$

In the first term in Eq.(A.28) only the zero-mode is projected out but in the second term higher modes contribute as well. Because the constraint equation (A.28) is nonlinear, the resulting quantum theory is as complicated as the formulation in usual coordinates. So far, it is not clear whether any dynamical simplifications (like "freezing out" of the zero-mode) arise in the infinite volume limit.

Bibliography

[Am 94] E. A. Ammons, Phys. Rev. **D50** (1994) 980.

[AS 70] M. Abramowitz and I. A. Stegun (Eds.), "Handbook of Mathematical Functions", (Dover, New York, 1970).

[BB 91] M. Burkardt and R. Busch, contributed to the Lake Louise Winter Institute: Particle Physics — The Factory Era, Lake Louise, Canada, February 17-23, 1991, (World Scientific, Singapore, 1991), eds.: B. A. Campbell, A. N. Kamal, P. Kitching and F. C. Khanna.

[BD 65] J. D. Bjorken and S. D. Drell: "Relativistic Quantum Fields", (Mc Graw Hill, New York, 1965).

[Pa 93] H.-C. Pauli, proceedings of Leipzig Workshop on Quantum Field Theory Theoretical Aspects of High Energy Physics, Bad Frankenhausen, Germany, 20-24 Sep 1993.

[BD+ 85] A. Bassetto, M. Dalbesco, I. Lazzizzera and R. Soldati, Phys. Rev. **D31** (1985) 2012; A. Basetto, G. Nardelli and R. Soldati, "Yang-Mills Theories in Algebraic Non-Covariant Gauges", (World Scientific, Singapore, 1991).

[BD+ 92] J. D. Bjorken, I. Dunietz and J. Taron, Nucl. Phys. **B371** (1992) 111.

[BD+ 94] A. Bassetto, F. De Biasio, L. Griguolo, Phys. Rev. Lett. **72** (1994) 3141.

[Be 56] P. G. Bergmann, Helv. Phys. Acta Suppl. **4** (1956) 79.

[Be 77] H. Bergknoff, Nucl. Phys. **B122** (1977) 215.

[BF+ 72] C. Buchiat, P. Fayet and N. Sourlas, Nuovo Cim. Lett. **4** (1972) 9.

[BG 78] I. Bars and M. B. Green, Phys. Rev. **D17** (1978) 537.

[BK+ 71] J. D. Bjorken, J. B. Kogut and D. E. Soper, Phys. Rev. **D3** (1971) 1382.

[BL 80] S. J. Brodsky and G. P. Lepage, Phys. Rev. **D22** (1980) 2157.

[BL 91a] M. Burkardt and A. Langnau, Phys. Rev. **D44** (1991) 1187.

[BL 91b] M. Burkardt and A. Langnau, Phys. Rev. **D44** (1991) 3857.

[BM+ 93] S. J. Brodsky, G. McCartor, H.-C. Pauli, S. S. Pinsky; Part. World **3**(1993) 109.

[BP 76] W. A. Bardeen and R. B. Pearson, Phys. Rev. **D14** (1976) 547; W. A. Bardeen, R. B. Pearson and E. Rabinovici, *ibid* **D21** (1980) 1037.

[BR+ 73] S. Brodsky, R. Roskies and R. Suaya, Phys. Rev. **D8** (1973) 4574.

[BP 91] S. J. Brodsky and H.-C. Pauli, Invited lectures given at 30th Schladming Winter School in Particle Physics: Field Theory, Schladming, Austria, Feb 27 - Mar 8, 1991.

[Bu 89a] M. Burkardt, Doktorarbeit, Erlangen, 1989.

[Bu 89b] M. Burkardt, Nucl. Phys. **A504** (1989) 762.

[Bu 89c] R. Busch, Diplomarbeit, Erlangen 1989.

[Bu 92a] M. Burkardt, Nucl. Phys. **B373** (1992) 371.

[Bu 92b] M. Burkardt, Nucl. Phys. **B373** (1992) 613.

[Bu 92c] M. Burkardt, in Proceedings to SPIN 92 (Nagoya, Japan, Nov 92).

[Bu 92d] M. Burkardt, Phys. Rev. **D46** (1992) R1924; R2751; M. Burkardt and E. Swanson, Phys. Rev. **D46** (1992) 5083.

[Bu 93] M. Burkardt, Phys. Rev. **D47** (1993) 4628.

[Bu 94a] M. Burkardt, Phys. Rev. **D49** (1994) 5446.

[Bu 94b] M. Burkardt, Workshop on "Quantum Infrared Physics", Paris, France, 1994, hep-ph/9409333.

[Bu 94c] M. Burkardt, proceedings to 'Theory of Hadrons and Light-Front QCD', Zakopane, August 1994, hep-ph/9410219.

[Bu 95] M. Burkardt, submitted to Phys. Rev. D., hep-ph/9505226.

[CC+ 76] C. G. Callan, N. Coote and D. J. Gross, Phys. Rev. **D13** (1976) 1649.

[CF+ 80] G. Curci, W. Furmanski and R. Petronzio, Nucl. Phys. **B175** (1980) 27.

[CL 84] T. P. Cheng and L. F. Li: "Gauge theory of elementary particle physics", (Oxford Univ. Press, Oxford, 1984).

[CM 69] S.-J. Chang and S. K. Ma, Phys. Rev. **180** (1969) 1506.

[Co 76] S. Coleman, Ann. Phys. (N.Y.) **101** (1976) 239.

[CR+ 73] S. Chang, R. Root and T. Yan, Phys. Rev. **D7** (1973) 1133.

[CY 72] S.-J. Chang and T.-M. Yan, Phys. Rev. **D7** (1972) 1147.

[CZ+ 95] C.-Y. Cheung, W.-M. Zhang and G.-L. Lin, hep-ph/9505232.

[Da 50] S. M. Dancoff, Phys. Rev. **78** (1950) 382.

[DH+ 75] R. F. Dashen, B. Hasslacher and A. Neveu, Phys. Rev. **D11** (1975) 3424; A. Zamolodchikov and A. Zamolodchikov, Ann. Phys. (N.Y.) **120** (1979) 253.

[Di 49] P. A. M. Dirac, Rev. Mod. Phys. **21** (1949) 392.

[Di 50] P. A. M. Dirac, Canad. J. Math. **1** (1950) 1; 'Lectures on Quantum Mechanics', (Benjamin, N.Y., 1964).

[DP 85] T. A. DeGrand and J. Potvin, Phys. Rev. **D31** (1985) 871.

[Ei 76] M. B. Einhorn, Phys. Rev. **D14** (1976) 3451; ibid. **D15** (1977) 1649, 3037.

[EP+ 87] T. Eller, H.-C. Pauli and S. J. Brodsky, Phys. Rev. **D35** (1987) 1493.

[EP 89] T. Eller and H.-C. Pauli, Z. Phys. **C42** (1989) 59.

[ES 95] M. Engelhardt and B. Schreiber, Z. Phys. **A 351** (1995) 71.

[FF 65] S. Fubini and G. Furlan, Physics **229** (1965).

[FJ 88] L. Fadeev and R. Jackiw, Phys. Rev. Lett. **60** (1988) 1962.

[FP 83] B. L. Friman, V. R. Pandharipande and R. B. Wiringa, Phys. Rev. Lett. **51** (1983) 763.

[GH+ 93] S. D. Glazek et al., Phys. Rev. **D47** (1993) 1599.

[GP 92] S. D. Glazek and R. J. Perry, Phys. Rev. **D45** (1992) 3740.

[Gr 92a] P. A. Griffin, Phys. Rev. **D46** (1992) 3538.

[Gr 92b] P. A. Griffin, Mod. Phys. Lett. **A7** (1992) 601; P. A. Griffin, Nucl. Phys. B **372** (1992) 270.

[Gr 93] P. A. Griffin, Phys. Rev. **D47** (1993) 1530.

[Gr 94a] P. Griffin, private communications.

[Gr 94b] P. Griffin, proceedings to 'Theory of Hadrons and Light-Front QCD', Zakopane, August 1994, hep-ph/9410243.

[GW 93] S. D. Glazek and K. G. Wilson, Phys. Rev. **D48** (1993) 5863; *ibid.* **D49** (1994) 4214.

[Ha 82] C. J. Hamer, Nucl. Phys. **B195** (1982) 503.

[Hi 91] J. R. Hiller, Phys. Rev. **D43** (1991) 2418; Phys. Rev. **D44** (1991) 2504; J. J. Wivoda and J. R. Hiller, Phys. Rev. **D47** (1993) 4647.

[HK+ 89] T. Heinzl, S. Krusche, E. Werner, Z. Phys. **A334** (1989) 443.

[HK+ 91a] T. Heinzl, S. Krusche, E. Werner, Phys. Lett. **B256** (1991) 55.

[HK+ 91b] T. Heinzl, S. Krusche, E. Werner, Phys. Lett. **B272** (1991) 54.

[HK+ 92a] T. Heinzl, S. Krusche, S. Simburger, E. Werner, Z. Phys. **C56** (1992) 415.

[HK+ 92b] T. Heinzl, S. Krusche, E. Werner, Phys. Lett. **B275** (1992) 410.

[HL 93] S. Huang and W. Lin, Ann. Phys. (N.Y.) **226** (1993) 248.

[HN+ 88] S. Huang, J. W. Negele and J. Polonyi, Nucl. Phys. **B307** (1988) 669.

[Ho 74] G.'t Hooft, Nucl. Phys. **B72** (1974) 461; ibid. **B75** (1974) 461.

[Ho 91] K. Hornbostel, in "From Fundamental Fields to Nuclear Phenomena", eds. J. A. McNeil and C. E. Price, (World Scientific, Singapore, 1991).

[Ho 92] K. Hornbostel, Phys. Rev. **D45** (1992) 3781.

[HP+ 90] K.Hornbostel, H.-C.Pauli and S.J.Brodsky, Proc. of the Ohio State Workshop on Relativistic Many-Body Physics, World Scientific, 1988;
S.J.Brodsky, Invited talk presented at the Third Lake Louise Winter Institute on QCD: Theory and Experiment, Chateau Lake Louise, Alberta, Canada, March 6-12, 1988;
K. Hornbostel, Ph.D.Thesis, SLAC-Report-333;
K. Hornbostel, S. J. Brodsky and H.-C. Pauli, Phys. Rev. **D41** (1990) 3814.

[HP 91] A. Harindranath and R. J. Perry, Phys. Rev. **D43** (1991) 492.

[HR 76] A. Hanson, T. Regge and C. Teitelboim, "Constrained Hamiltonian Systems", Academia Nazionale dei Lincei, (Rome, 1976).

[HS+ 82] J. E. Hirsch, R. L. Sugar, D. J. Scalapino and R. Blankenbecler, Phys. Rev. **B26** (1982) 5033; J. Potvin and T. A. DeGrand, Phys. Rev. **D30** (1984) 1285.

[HV 87] A. Harindranath and J. P. Vary, Phys. Rev. **D36** (1987) 1141; **D37** (1988) 1064; 1076; 3010;
W.-M. Zhang, Chin. J. Phys. **32** (1994) 717. C. J. Benesh and J. P. Vary, Z. Phys. **C49** (1991) 411.

[HZ 93] A. Harindranath and W.-M. Zhang, Phys. Rev. **D 48** (1993) 4868; 4881; 4903.

[Ja 72] R. Jackiw, *Springer Tracts in Modern Physics* **62**, ed. G. Höhler (Springer, 1972).

[Ja 85] R. L. Jaffe, in "Los Alamos School on Relativistic Dynamics and Quark-Nuclear Physics", ed. M. B. Jackson and A. Picklesimer, (Wiley, New York, 1985).

[Ji 93] X. Ji, Comments Nucl. Part. Phys. **21** (1993) 123.

[KP 92] M. Kaluža and H.-C. Pauli, Phys. Rev. **D45** (1992) 2968;
M. Krautgärtner, H.-C. Pauli and F. Wölz, Phys. Rev. **D45** (1992) 3755;
L. C. L. Hollenberg, K. Higashijima, R. C. Warner and B. H. J. McKellar, Prog. Theor. Phys. **87** (1992) 441.

[KP 93] A. C. Kalloniatis and H. C. Pauli,Z. Phys. **C60** (1993) 255; Z.Phys. **C63** (1994) 161;
A. C. Kalloniatis and D. G. Robertson, Phys. Rev. **D50** (1994) 5262.

[KP+ 94] A. C. Kalloniatis, H.-C. Pauli and S. S. Pinsky, Phys. Rev. **D50** (1994) 6633.

[KS 70] J. B. Kogut and D. E. Soper, Phys. Rev. **D1** (1970) 2901.

[LB 93] A. Langnau and M. Burkardt, Phys. Rev. **D47** (1993) 3452.

[Li 86] M. Li, Phys.Rev. **D34** (1986) 3888;
M. Li, L. Wilets and M. C. Birse, J.Phys. **G13** (1987) 915.

[LN+ 94a] F. Lenz, H. W. L. Naus, K. Ohta and M. Thies, Ann. Phys. (N.Y.) **233** (1994) 17.

[LN+ 94b] F. Lenz, H. W. L. Naus, K. Ohta and M. Thies, Ann. Phys. (N.Y.) **233** (1994) 51.

[LN+ 94c] F. Lenz, H. W. L. Naus and M. Thies, Ann. Phys. (N.Y.) **233** (1994) 317.

[LS 93] H.-H. Lin and D. Soper, Phys. Rev. **D48** (1993) 1841.

[LT 89] F.Lenz and M.Thies, private communications.

[LT 91] F. Lenz, S. Levit, M. Thies and K. Yazaki, Ann. Phys.(N.Y.) **208** (1991) 1; F. Lenz, Proceedings of NATO Advanced Study Institute on "Hadrons and Hadronic Matter", eds. Vautherin et al. (Plenum, New York, 1990).

[Ma 83] S. Mandelstam, Nucl. Phys. **B213** (1983) 149;
G. Leibbrandt, Phys. Rev. **D29** (1984) 1699; Rev. Mod. Phys. **59** (1987) 1067.

[Ma 85] N. S. Manton, Ann. Phys. (N.Y.) **159** (1985) 220.

[Ma 92] K. Mahrt, Diplomarbeit, Erlangen, 1992.

[Mc 88] G. McCartor, Z. Phys. **C41** (1988) 271; **C52** (1991) 611; **C64** (1994) 349.

[MP+ 91] D. Mustaki, S. Pinsky, J. Shigemitsu and K. G. Wilson, Phys. Rev. **D43** (1991) 3411.

[Mu 88] D. Mustaki, Phys. Rev. **D38** (1988) 1260.

[MV 94] L. McLerran and R. Venugopalan, Phys. Rev. **D49** (1994) 2233; 3352; *ibid.* **D50** (1994) 2225.

[NO 87] J. W. Negele and H. Orland, "Quantum Many-Particle Systems", (Addison Wesley, Redwood City, 1987).

[NS+ 81] V. A. Novikov, M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. **B191** (1981) 301.

[Pa 93] H.-C. Pauli, proceedings of Leipzig Workshop on Quantum Field Theory Theoretical Aspects of High Energy Physics, Bad Frankenhausen, Germany, 20-24 Sep 1993.

[PB 85] H.-C. Pauli and S. J. Brodsky, Phys. Rev. **D32** (1985) 1993, 2001.

[Pe 94a] R. J. Perry, Ann. Phys. (N.Y.) **232** (1994) 116.

[Pe 94b] R. J. Perry, invited lectures presented at 'Hadrons 94', Gramado, Brasil, April, 1994.

[PF 89] E. V. Prokhvatilov and V. A. Franke, Sov. J. Nucl. Phys. **49** (1989) 688.

[PH+ 90] R. J. Perry, A. Harindranath and K. G. Wilson, Phys. Rev. Lett. **65** (1990) 2959.

[PH 91] R. J. Perry and A. Harindranath, Phys. Rev. **D43** (1991) 4051.

[PN+ 95] E. V. Prokhvatilov, H.W.L. Naus and H.-J. Pirner, Phys. Rev. **D51** (1995) 2933.

[PV 93] S. S. Pinsky and B. Van de Sande, Phys. Rev. **D48** (1993) 816.

[PV 94] S. S. Pinsky and B. Van de Sande, Phys. Rev. **D49** (1994) 2001.

[PV+ 95] S. S. Pinsky, B. Van de Sande and J. R. Hiller, to appear in Phys. Rev. D

[PW 93] R. J. Perry and K. G. Wilson, Nucl. Phys. **B403** (1993) 403.

[RM 92] D. G. Robertson and G. McCartor, Z. Phys. **C53** (1992) 661; G. McCartor and D. G. Robertson, Z. Phys. **C53** (1992) 679.

[RM 94] G. McCartor and D. G. Robertson, Z. Phys.**C62** (1994) 349; hep-ph/9501107, to appear in Z. Phys. **C**.

[Ro 70] F. Röhrlich, Acta Phys. Aust. **32** (1970) 87.

[Ro 93] D. G. Robertson, Phys. Rev. **D47** (1993) 2549.

[St 92] S. Stampfer, Diplomarbeit, Erlangen, 1992.

[Su 68] L. Susskind, Phys. Rev. **165** (1968) 1535.

[Su 82] K. Sundermayer, 'Constrained Dynamics', Lecture Notes in Physics 169, (Springer, Berlin, 1982).

[Ta 45] I. Tamm, J. Phys. (USSR) **9** (1945) 449.

[TB 91] A. C. Tang, S. J. Brodsky and H.-C. Pauli, Phys. Rev. **D44** (1991) 1842.

[Th 79] C. B. Thorn, Phys. Rev. **D19** (1979) 639; *ibid* **D20** (1979) 1934.

[VF 94] J. Vary and T. J. Fields, proceedings to 'Theory of Hadrons and Light-Front QCD', Zakopane, August 1994, hep-ph/9411263.

[We 69] S. Weinberg, Phys. Rev. **150** (1966) 1313.

[Wi 75] K. G. Wilson, Rev. Mod. Phys. **47** (1975) 773.

[Wi 91] K. G. Wilson, talk at Aspen Center for Physics, Aspen, CO, 1991 (a video tape of this lecture is available from S. Pinsky).

[Wo 93] P. M. Wort, Phys. Rev. **D47** (1993) 608.

[WR 94] K. G. Wilson and D. G. Robertson, proceedings to 'Theory of Hadrons and Light-Front QCD', Zakopane, August 1994, hep-ph/9411007.

[Wu 77] T. T. Wu, Phys. Lett. **61B** (1977) 142; Phys. Rep. **49** (1979) 245.

[WW+ 94] K. G. Wilson et al., Phys. Rev. **D49** (1994) 6720.

[Yn 83] F. J. Yndurain: "Quantum Chromodynamics", (Springer, New York, 1983).

[Zh 85] A. R. Zhitnitsky, Phys. Lett. **165B** (1985) 405; Sov. J. Nucl. Phys. **43** (1986) 999; Sov. J. Nucl. Phys. **44** (1986) 139.

This figure "fig1-1.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9505259v1

This figure "fig2-1.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9505259v1

This figure "fig3-1.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9505259v1

This figure "fig4-1.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9505259v1

This figure "fig1-2.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9505259v1

This figure "fig2-2.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9505259v1

This figure "fig3-2.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9505259v1

This figure "fig4-2.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9505259v1

This figure "fig1-3.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9505259v1

This figure "fig2-3.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9505259v1

This figure "fig3-3.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9505259v1

This figure "fig4-3.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9505259v1

This figure "fig4-4.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9505259v1